

# Recent star formation in high-redshift early-type galaxies: insights from the rest-frame UV

S. Kaviraj<sup>1</sup>†, S. K. Yi<sup>2</sup>, E. Gawiser<sup>3</sup>, P. G. van Dokkum<sup>3</sup>,  
S. Khochfar<sup>1</sup>, K. Schawinski<sup>1</sup> and J. Silk<sup>1</sup>

<sup>1</sup>Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, U.K.

<sup>2</sup>Yonsei University, Centre for Space Astrophysics, Seoul 120749, Korea

<sup>3</sup>Yale Center for Astronomy and Astrophysics, Yale University, New Haven, CT 06520-8121, U.S.A.

**Abstract.** We combine deep *UBVRIzJK* photometry from the MUSYC survey with redshifts from the COMBO-17 survey to study the *rest-frame* ultraviolet (*UV*) properties of 674 high-redshift ( $0.5 < z < 1$ ) early-type galaxies, drawn from the Extended Chandra Deep Field South (E-CDFS). Galaxy morphologies are determined through visual inspection of Hubble Space Telescope (HST) images taken from the GEMS survey. We harness the sensitivity of the *UV* to young ( $< 1$  Gyrs old) stars to quantify the *recent* star formation history of the early-type population. We find compelling evidence that early-types of all luminosities form stars over the lifetime of the Universe, although the bulk of their star formation is already complete at high redshift. Luminous ( $-23 < M(V) < -20.5$ ) early-types form 10-15 percent of their mass after  $z = 1$ , while their less luminous ( $M(V) > -20.5$ ) counterparts form 30-60 percent of their mass in the same redshift range.

**Keywords.** galaxies: elliptical and lenticular, cD, galaxies: evolution, galaxies: formation, galaxies: stellar contents, ultraviolet: galaxies


---

## 1. Introduction

The vast majority of work on early-type galaxies in the past has focussed on optical spectro-photometric data. A significant drawback of optical photometry is its lack of sensitivity to moderate amounts of recent star formation (RSF). While red optical colours imply a high-redshift formation epoch for the *bulk* of the stellar population in early-type galaxies, the optical spectrum remains largely unaffected by the minority of stellar mass that forms in these systems at low and intermediate redshift. As a result it is virtually impossible to quantify early-type star formation histories (SFHs) over the last 8 billion years ( $0 < z < 1$ ).

RSF can be efficiently traced using the rest-frame ultraviolet (*UV*) spectrum, which is sensitive to young, massive main sequence stars with ages less than  $\sim 1$  Gyr. Using a large sample of early-type galaxies detected by the GALEX *UV* space telescope, Kaviraj *et al.* (2006) have recently shown that, contrary to the expectations of traditional ‘monolithic collapse’ models (e.g. Bower *et al.* 1992), local early-types show widespread recent star formation (RSF) - at least 30 percent show blue *UV*-optical colours indicative of unambiguous RSF in these systems. The work presented here extends these results to high redshift by exploiting deep optical photometry to trace the rest-frame *UV* properties of early-type galaxies in the redshift range  $0.5 < z < 1$ .

† skaviraj@astro.ox.ac.uk



**Figure 1.** LEFT-HAND PANEL: Rest-frame ( $NUV - r$ ) colour-magnitude relation of the E-CDFS galaxy population. RIGHT-HAND PANEL: Rest-frame ( $NUV - r$ ) colour plotted against the RSF in the E-CDFS galaxy population. The  $NUV$  ( $\sim 2300\text{\AA}$ ) and  $r$  band filters are taken from the GALEX and SDSS filtersets respectively.

## 2. Estimation of recent star formation

The RSF in individual galaxies is calculated by comparing their multi-wavelength photometry with synthetic galaxy populations, generated in the framework of the standard model. Synthetic populations are generated using the semi-analytical model of Khochfar and Burkert (2003). A library of synthetic photometry is constructed by combining each of  $\sim 15,000$  model galaxies with a single metallicity in the range  $0.1Z_{\odot}$  to  $2.5Z_{\odot}$ , dust extinction parametrised by a value of  $E(B - V)$  in the range 0 to 0.5 and convolving them with the stellar models of Yi (2003). We construct an ‘average SFH’ for each galaxy, by combining the SFHs of all models in the library weighted by their individual likelihoods. The RSF, defined as the *mass fraction of stars formed within the last Gyr*, is calculated from this average SFH. Figure 1 shows the derived rest-frame ( $NUV - r$ ) colour-magnitude relation and the values of RSF in the E-CDFS galaxy population.

## 3. Summary and discussion

The early-type population as a whole exhibits a typical RSF between 5 and 13 percent in the redshift range  $0.5 < z < 1$ , while the early-types on the broad ‘red sequence’, ( $NUV - r > 4$ ), typically show RSF values less than 5 percent. The reddest early-types (which are also the most luminous) are virtually quiescent with RSF values of  $\sim 1$  percent. Since the timescale of this study is  $\sim 2.5$  Gyrs, a simple extrapolation (from RSF values in Figure 1) indicates that luminous ( $-23 < M(V) < -20.5$ ) early-type galaxies typically form up to 10-15 percent of their mass after  $z = 1$  (with a tail to higher values), while less luminous early-types ( $M(V) > -20.5$ ) form 30-60 percent of their mass after  $z = 1$ .

## References

- Bower, R. G., Lucey, J. R., & Ellis, R. 1992, *MNRAS* 254, 589  
 Kaviraj, S. & GALEX Science Team 2006, *ApJ GALEX dedicated issue, Dec 2007*  
 Khochfar, S. & Burkert, A. 2003, *ApJ* 597, L117  
 Yi, S. K. 2003, *ApJ* 582, 202



**Figure 1.** LOC member Marc Sarzi and Jesús Falcón-Barroso in the lecture theatre.




**Figure 2.** George Bendo and Lisa Young discussing during a coffee break. LOC member Davor Krajnović is in the centre background.



**Figure 3.** Elena D'Onghia and Eric Emsellem discussing during a coffee break. LOC member Davor Krajnović and Robert Zylka are visible in the left background, John Kormendy and SOC member Matthias Steinmetz on the right.



**Figure 4.** Thorsten Naab, Françoise Combes, Peter Erwin and Eric Emsellem discussing during a coffee break.



**Figure 5.** Anatoly Klypin and Daniel Ceverino during the symposium dinner.



**Figure 6.** SOC member David Koo, Ryan Houghton and Mark Ammons discussing during the symposium dinner. Giuseppe Aronica is in the background.





**Figure 7.** John Kormendy, Elena Dalla Bonta and Enrico Maria Corsini discussing during a coffee break. Lauren MacArthur is visible in the background on the right.



**Figure 8.** Brad Gibson, Rosa Domínguez-Tenreiro, Daniel Ceverino, Andres Escala and Anatoly Klypin discussing during a coffee break.