

INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, <http://www.editorialmanager.com/psm/>. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

1. S.I. units should be used throughout in text, figures and tables.
2. Authors should spell out in full any abbreviations used in their manuscripts.
3. Foreign quotations and phrases should be followed by a translation.
4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* **286**, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* **32**, 173–179.

Cleckley HJ (1941). *The Mask of Sanity*, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank.org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq_tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. **Line artwork:** Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; **Combination artwork (line/tone):** Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; **Black and white halftone artwork:** Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; **Colour halftone artwork:** Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. *Tables* Tables should be numbered consecutively in the text in Arabic numerals and each typed on a separate sheet after the References section. Titles should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLE

The association between cognitive function and subsequent depression: a systematic review and meta-analysis

Scult MA, Paulli AR, Mazure ES, Moffitt TE, Hariri AR & Strauman TJ 1

COMMENTARY

The DSM revision process: needing to keep an eye on the empirical ball

First MB 19

ORIGINAL ARTICLES

Childhood interleukin-6, C-reactive protein and atopic disorders as risk factors for hypomanic symptoms in young adulthood: a longitudinal birth cohort study

Hayes JF, Khandaker GM, Anderson J, Mackay D, Zammit S, Lewis G, Smith DJ & Osborn DPJ 23

Emotional intelligence and non-social cognition in schizophrenia and bipolar I disorder

Frajo-Apor B, Kemmler G, Pardeller S, Plass T, Mühlbacher M, Welte A-S, Fleischhacker WW & Hofer A 35

Cognitive-behaviour therapy for post-traumatic stress in schizophrenia. A randomized controlled trial

Steel C, Hardy A, Smith B, Wykes T, Rose S, Enright S, Hardcastle M, Landau S, Baksh MF, Gottlieb JD, Rose D & Mueser KT 43

Cost-effectiveness of cognitive remediation and supported employment for people with mental illness: a randomized controlled trial

Yamaguchi S, Sato S, Horio N, Yoshida K, Shimodaira M, Taneda A, Ikebuchi E, Nishio M & Ito J 53

Effects of a transdiagnostic unguided Internet intervention ('velibra') for anxiety disorders in primary care: results of a randomized controlled trial

Berger T, Urech A, Krieger T, Stolz T, Schulz A, Vincent A, Moser CT, Moritz S & Meyer B 67

Comparative risk of self-harm hospitalization amongst depressive disorder patients using different antidepressants: a population-based cohort study in Taiwan

Wu C-S, Liao S-C, Tsai Y-T, Chang S-S & Tsai H-J 81

A study of wrist-worn activity measurement as a potential real-world biomarker for late-life depression

O'Brien JT, Gallagher P, Stow D, Hammerla N, Ploetz T, Firbank M, Ladha C, Ladha K, Jackson D, McNaney R, Ferrier IN & Olivier P 93

Religiosity, race/ethnicity, and alcohol use behaviors in the United States

Meyers JL, Brown Q, Grant BF & Hasin D 103

The longitudinal relationship between post-traumatic stress disorder and perceived social support in survivors of traumatic injury

Nickerson A, Creamer M, Forbes D, McFarlane AC, O'Donnell ML, Silove D, Steel Z, Felmingham K, Hadzi-Pavlovic D & Bryant RA 115

Childhood bullying victimization is associated with use of mental health services over five decades: a longitudinal nationally representative cohort study

Evans-Lacko S, Takizawa R, Brimblecombe N, King D, Knapp M, Maughan B & Arseneault L 127

Interrelationship between family history of alcoholism and generational status in the prediction of alcohol dependence in US Hispanics

Chartier KG, Thomas NS & Kendler KS 137

Symptoms of post-traumatic stress disorder, severe psychological distress, explosive anger and grief amongst partners of survivors of high levels of trauma in post-conflict Timor-Leste

Silove DM, Tay AK, Steel Z, Tam N, Soares Z, Soares C, dos Reis N, Alves A & Rees S 149

The impact of postpartum post-traumatic stress disorder symptoms on child development: a population-based, 2-year follow-up study

Garthus-Niegel S, Ayers S, Martini J, von Soest T & Eberhard-Gran M 161

Effects of early life stress on depression, cognitive performance and brain morphology

Saleh A, Potter GG, McQuoid DR, Boyd B, Turner R, MacFall JR & Taylor WD 171

Correspondence 183-187