

### (Q86) Vulnerable Populations: A Social Ecological Approach to Nursing Research of Chronic Illness in Disaster Management

Joan Valas

University of Pennsylvania School of Nursing, Park Ridge, New Jersey USA

**Statement of Purpose:** This presentation describes the need for a social ecological approach to nursing research of disasters and the impact on the health of vulnerable populations with chronic illnesses. Such a framework ranges from individual, social networks, community, and societal dimensions. Chronic illnesses and disasters both have multi-factor etiologies and their complexities merit this broad approach. This presentation examines the social forces that predispose groups of people to be more vulnerable than others.

**Learning Objectives:** The learner will be able to: (1) describe the social forces that predispose groups of people with chronic illnesses to be more vulnerable than others to the impact of disasters; (2) describe interdependent dimensions of the social ecological model at which disasters and determinants of health may be understood broadly; and (3) recognize the impact of this approach to nursing research in addressing the need of vulnerable populations with chronic illnesses during a disaster.

**Keywords:** chronic illness; disaster management; emergency health; nursing; research; vulnerable populations

*Prehosp Disast Med* 2009;24(2):s9

### (Q87) Investigation of the Frequency of Different Emergency Calls in Northern and Southern Geographical Areas of Tehran

Mohamad Kalantari Meibodi; Afshin Amini

Shahid Beheshti University, Tehran, Iran

**Introduction:** Urban society hosts many different cultural, economic, and social factors that may be the cause of specific emergency calls. Because the process of decision-making should be based on necessity, it is important to gather and organize information from the 115 emergency centers in each area and identify the kinds of medical expertise required to cope with the health-related problems of that area.

**Methods:** Four emergency teams, each consisting of two trained technicians, were equipped with one vehicle. Two of the teams were deployed to southeastern Tehran while the other two were deployed to the northeast. These teams were commissioned to transfer patients to the closest hospital after receiving emergency calls. After assigning patients into categories of cardiovascular, bronchitis, non-drug abuse suicidal, poisoning, and accidents, the following results were observed.

**Results:** Car crashes were the most frequent causes of emergency calls both in southern and in northern areas of Tehran. The number of accidents and the related frequency of death and injury were greater for northern Tehran and the causes for most of these cases were accidents. The only cause of emergency calls for which there were more women than men was, suicide. In both the

north and south areas, car accidents and poisonings happened mostly to those between 18 and 27 years of age. In northern areas of Tehran, addictive drug abuse was more frequent among men, while non-addictive drug abuse was more frequent among women.

**Keywords:** emergency calls; Iran; public health; social

*Prehosp Disast Med* 2009;24(2):s9

### (Q88) Mortality Rates for Pedestrian-Motor Vehicle Collisions in Mexico City, 2004–2006

Oscar Santiago-Espinosa; Martha Hajar-Medina

Instituto Mexicano Del Seguro Social, Cancun, Mexico

**Objective:** To determine the factors related to the mortalities caused for pedestrian-motor vehicle collisions in Mexico City from 2004–2006.

**Methods:** Information was gathered from death certificates of DGE SS. When the cause was recorded under the code “ICD X: V01-V09”, the INEGI was used. The net, specific, adjusted, and registered rates were calculated. The genre, age, residence, and the accident site and time were analyzed.

**Results:** Of 2,119 deaths, 68% (1,452) were residents of Mexico City. The average specified fatality was 2.5 (CI 95% 2.20–2.75). Men had a higher risk than women. The mortality rate increased with age; those 74 years of age had a rate of 185.3 for men and 75.6 for women/per one thousand population. Of the deaths, 65% occurred in the 15–64-years-old age group. The net rate was 16.6/one thousand from a general population. The districts with the highest adjusted rates were Cuauhtemoc, Xochimilco, and Venustiano Carranza with 22.6, 20.4, and 20.3/one thousand, respectively. The internal and external rates indicate that the districts of Gustavo A. Madero and Miguel Hidalgo have twice the risk of death in the same district. The fatalities occurred on Friday, Saturday, and Sunday, making a total of 50%. Only 749 (35.3%) recorded the accident site. Iztapalapa and Gustavo A. Madero are notable with 19% and 16% of the total fatalities, respectively.

**Conclusions:** The high mortality rate of people at a productive age impacts the stability and integrity of the families. Hardly any data are recorded on accident sites. Therefore, it is necessary to improve the information-gathering systems with specific means of measuring prevention.

**Keywords:** accident sites; Mexico City; mortality rates; pedestrian-motor vehicle collisions; public health

*Prehosp Disast Med* 2009;24(2):s9

### (Q89) Missed Opportunities: HIV Testing in a Bronx, New York Emergency Department

Andrew Park

St. Barnabas Hospital, New York, New York USA

**Background:** Undiagnosed HIV infection is a serious and growing problem in the world, but particularly in places like the South Bronx in New York City where a vulnerable, primarily African-American and Latino population, suffers some of the highest rates of newly diagnosed HIV infections in the United States.

**Objective:** In this study, the lack of HIV testing and counseling in a South Bronx emergency department is addressed by demonstrating the frequency of high-risk