THE PREPARATION OF MANUSCRIPTS

The attention of authors is particularly directed to the following requests.

1. Papers should be typed, double-spaced, on one side of white paper (of which A4, 210 by 297 mm, is a suitable size). The pages must be numbered. Margins of 30 mm should be left at the side, top and bottom of each page. Two clear copies should be sent.

A cover page should give the title, the author's name and institution, with the address at which mail is

to be sent.

The title, while brief, must be informative (e.g. A new proof of the prime-number theorem, whereas Some applications of a theorem of G. H. Hardy would be useless).

The first paragraph or two should form a summary of the main theme of the paper, providing an

abstract intelligible to mathematicians.

For a typescript to be accepted for publication, it must accord with the standard requirements of publishers, and be presented in a form in which the author's intentions regarding symbols etc. are clear to a printer (who is not a mathematician).

The following notes are intended to help the author in preparing the typescript. New authors may well enlist the help of senior colleagues, both as to the substance of their work and the details of setting it out correctly and attractively.

2. Notation

Notation should be chosen carefully so that mathematical operations are expressed with all possible neatness, to lighten the task of the compositor and to reduce the chance of error.

For instance n_k (n sub k) is common usage, but avoid if possible using c sub n sub k. Fractions are generally best expressed by a solidus. Complicated exponentials like

$$\exp\{z^2\sin\theta/(1+y^2)\}$$

should be shown in this and no other way.

In the manuscript, italics, small capitals and capitals are specified by single, double and triple underlinings. Bold faced type is shown by wavy underlining; wavy will be printed wavy.

It helps if displayed equations or statements which will be quoted later are numbered in order on

the right of their line. They can then be referred to by, for example, 'from (7)'.

The author must enable the printer (if necessary by pencilled notes in the margin) to distinguish between similar symbols such as o, O o, O, O; x, X, \times ; ϕ , Φ , \emptyset ; I, I; ε , \in ; κ , k.

Greek letters can be denoted by Gk in the margin.

If an author wishes to mark the end of the proof of a theorem, the sign I may be used. Footnotes should be avoided.

3. Diagrams

It is extremely helpful if diagrams are drawn in Indian ink on white card, faintly blue or green-lined graph paper, or tracing cloth or paper. Symbols, legends and captions should be given on a transparent overlay. Each text figure must be numbered as Figure 1, Figure 2, ... and its intended position clearly indicated in the manuscript:

Figure 1 here

The author's name in pencil must be on all separate sheets of diagrams.

A figure is expensive to reproduce and should be included only when the subject matter demands it,

or when it greatly clarifies the exposition.

The Society recognizes that some authors do not have the facilities for producing drawings of a sufficiently high standard to be reproduced directly and it is therefore willing to have such diagrams redrawn, provided that they are clear.

4. Tables

Tables should be numbered (above the table) and set out on separate sheets. Indicate the position of each in the text as for figures:

Table 3 here

5. References

References should be collected at the end of the paper numbered in alphabetical order of the authors' names. Titles of journals should be abbreviated as in *Mathematical Reviews*. The following examples show the preferred style for references to a paper in a journal, a paper in a proceedings volume, a book and an unpublished dissertation:

[1] J. F. Adams. On the non-existence of elements of Hopf invariant one. Ann. of Math. (2) 72

(1960), 20-104

[2] M. P. FOURMAN and D. S. Scott. Sheaves and logic. In Applications of Sheaves, Lecture Notes in Math. vol. 753 (Springer-Verlag, 1979), pp. 302-401.

[3] P. T. JOHNSTONE. Stone Spaces. Cambridge Studies in Advanced Math. no. 3 (Cambridge University Press, 1982).

[4] F. W. LAWVERE. Functorial semantics of algebraic theories. Ph.D. thesis. Columbia University (1963).

Mathematical Proceedings of the Cambridge Philosophical Society

MPCPCO 114 (Pt 2) 191-382 (1993) 0305-0041 September 1993

CONTENTS

	PAGE
Bennett, M. Fractional parts of powers of rational numbers	. 191
MERRIMAN, J. R. & SMART, N. P. Curves of genus 2 with good reduction away from 2 with a	
rational Weierstrass point	. 203
HILLMAN, J. A. On 3-dimensional Poincaré duality complexes and 2-knot groups	. 215
Kashiwabara, T. Mod p K-theory of $\Omega^{\infty}\Sigma^{\infty}X$ revisited	. 219
Promislow, S. D. A class of groups producing simple, unique trace C*-algebras	. 223
MELBOURNE, I. & DELLNITZ, M. Normal forms for linear Hamiltonian vector fields commuting	
with the action of a compact Lie group	235
Hall, J. I. The general theory of 3-transposition groups.	269
Gupta, N. & Shpilrain, V. Nielsen's commutator test for two-generator groups	295
FOUNTAIN, J. & LEWIN, A. Products of idempotent endomorphisms of an independence algebra of	
infinite rank	303
Bowers, P. L. The upper Perron method for labelled complexes with applications to circle	-
packings	321
Arratia, R., Barbour, A. D. and Tavaré, S. On random polynomials over finite fields	347
Rogers, L. C. G. The harmonic functions of $(A_l, B_l)^1$	369
Kondracki, W., Królak, A. & Rudnicki, W. On the instability of a class of generalized strong	
	379
curvature singularities	319

© The Cambridge Philosophical Society 1993 Printed in Great Britain by the University Press, Cambridge

CAMBRIDGEUNIVERSITY PRESS

