

ON PURELY PERIODIC BETA-EXPANSIONS OF PISOT NUMBERS

YUKI SANO

Abstract. We characterize numbers having purely periodic β -expansions where β is a Pisot number satisfying a certain irreducible polynomial. The main tool of the proof is to construct a natural extension on a d -dimensional domain with a fractal boundary.

§1. Introduction

Let $\beta > 1$ be a real number and let T_β be the β -transformation on the unit interval $[0, 1)$ given by

$$T_\beta x = \beta x - [\beta x],$$

where $[x]$ denotes the integer part of x . Then every $x \in [0, 1)$ can be written as

$$x = \sum_{k=1}^{\infty} b_k \beta^{-k}, \quad b_k = [\beta T_\beta^{k-1} x].$$

We call this representation in base β the β -expansion, which was introduced by Rényi [16]. It is denoted by

$$x = .b_1 b_2 \dots$$

A real number $x \in [0, 1)$ is said to have an eventually periodic β -expansion with period p if there exist integers $m \geq 0$ and $p \geq 1$ such that

$$x = .b_1 b_2 \dots b_m (b_{m+1} b_{m+2} \dots b_{m+p})^\infty,$$

where w^∞ will denote the sequence $www \dots$. In particular, if we can choose $m = 0$, we say that x has a purely periodic β -expansion with period p , that is,

$$x = .(b_1 b_2 \dots b_p)^\infty.$$

Received February 1, 2000.

1991 Mathematics Subject Classification: 34C35, 58F22.

We know that x has a purely periodic β -expansion with period p if and only if $T_\beta^p x = x$.

For $x = 1$, we can define the β -expansion of 1 in the same way:

$$d(1, \beta) = .t_1 t_2 \dots, \quad t_k = \lfloor \beta T_\beta^{k-1} 1 \rfloor.$$

Let D_β be the set of β -expansions of numbers in $[0, 1)$. Parry characterized the set D_β in [13]. By $<_{lex}$ will be denoted the lexicographical order, that is, $(v_i)_{i=1}^\infty <_{lex} (w_i)_{i=1}^\infty$ means that there exists $k \geq 1$ such that $v_j = w_j$ for any $1 \leq j < k$ and $v_k < w_k$. The (one-sided) shift σ_s maps a point $(v_i)_{i=1}^\infty$ to the point $(v'_i)_{i=1}^\infty = \sigma_s((v_i)_{i=1}^\infty)$ whose i th coordinate is given by $v'_i = v_{i+1}$.

THEOREM (PARRY). *Let $\beta > 1$ be a real number, and let $d(1, \beta) = .t_1 t_2 \dots$. Let w be an infinite sequence of positive integers.*

(1) *If $d(1, \beta)$ is infinite,*

$$w \in D_\beta \iff \forall u \geq 0, \sigma_s^u(w) <_{lex} d(1, \beta).$$

(2) *If $d(1, \beta)$ is finite, $d(1, \beta) = .t_1 \dots t_{n-1} t_n$, say, then*

$$w \in D_\beta \iff \forall u \geq 0, \sigma_s^u(w) <_{lex} d^*(1, \beta) = (t_1 \dots t_{n-1} (t_n - 1))^\infty.$$

Bertrand [3] and K. Schmidt [18] investigated eventually periodic β -expansions. A Pisot number is an algebraic integer (> 1) whose conjugates other than itself have modulus less than one. Let $\mathbb{Q}(\beta)$ be the smallest extension field of rational numbers \mathbb{Q} containing β .

THEOREM (BERTRAND, K. SCHMIDT). *Let β be a Pisot number and let x be a real number in $[0, 1)$. Then x has an eventually periodic β -expansion if and only if $x \in \mathbb{Q}(\beta)$.*

In [1], Akiyama gives a sufficient condition for pure periodicity where β belongs to a certain class of Pisot numbers. Hara and Ito characterized purely periodic modified β -expansions for a quadratic irrational number β in [8]. The present author studied necessary and sufficient condition for pure periodicity in [11] where β is a cubic Pisot number whose minimal polynomial is given by

$$Irr(\beta) = x^3 - k_1 x^2 - k_2 x - 1, \quad k_1 (\neq 0), \quad k_2 \in \mathbb{N} \cup \{0\}, \quad \text{and } k_1 \geq k_2.$$




Figure 1: Figure of \widehat{Y} in case $d = 3$.

In this paper, we will generalize the results of [11]. Hereafter, β is a positive root of the polynomial:

$$\begin{aligned} Irr(\beta) &= x^d - k_1x^{d-1} - k_2x^{d-2} - \dots - k_{d-1}x - 1, \\ k_i &\in \mathbb{Z}, \text{ and } k_1 \geq k_2 \geq \dots \geq k_{d-1} \geq 1. \end{aligned}$$

Then β is a Pisot number. We have the following result:

MAIN THEOREM. *Let x be a real number in $\mathbb{Q}(\beta) \cap [0, 1)$. Then x has a purely periodic β -expansion if and only if x is reduced.*

We define reduced numbers in Section 5. For our purpose, we introduce a d -dimensional domain \widehat{Y} with a fractal boundary (see Figure 1 and the definition in Section 4) and a natural extension of T_β on \widehat{Y} , which were originally discussed in [14] and [19]. In [8] and [9], you can find the basic idea of the proof.

I would like to thank Professor S. Akiyama and Professor Sh. Ito for many helpful suggestions and encouragement. I am grateful to N. Tangiku for a life filled with love and support. Finally, I am greatly grateful to the referee for careful reading the manuscript and giving helpful comments.

§2. Admissible sequences of β -expansions

Recall that β is a positive root of the irreducible polynomial

$$\begin{aligned} (2.1) \quad Irr(\beta) &= x^d - k_1x^{d-1} - k_2x^{d-2} - \dots - k_{d-1}x - 1, \\ k_i &\in \mathbb{Z}, \text{ and } k_1 \geq k_2 \geq \dots \geq k_{d-1} \geq 1. \end{aligned}$$

From [4], we know that β is a Pisot number. From Theorem (Parry) it follows that

$$d(1, \beta) = .k_1 k_2 \dots k_{d-1} 1.$$

Let

$$\beta = \beta^{(1)}, \beta^{(2)}, \dots, \beta^{(r_1)}$$

be the real Galois conjugates and

$$\beta^{(r_1+1)}, \overline{\beta^{(r_1+1)}}, \beta^{(r_1+2)}, \overline{\beta^{(r_1+2)}}, \dots, \beta^{(r_1+r_2)}, \overline{\beta^{(r_1+r_2)}}$$

be the complex Galois conjugates of β , where $r_1 + 2r_2 = d$ and \bar{v} is the complex conjugate of a complex number v . The corresponding conjugates of $x \in \mathbb{Q}(\beta)$ are also denoted by

$$x = x^{(1)}, \dots, x^{(r_1)}, x^{(r_1+1)}, \overline{x^{(r_1+1)}}, \dots, x^{(r_1+r_2)}, \overline{x^{(r_1+r_2)}}.$$

Let M be the companion matrix of the polynomial (2.1), that is,

$$M = \begin{bmatrix} k_1 & k_2 & \dots & k_{d-1} & 1 \\ 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 0 \end{bmatrix}.$$

We know that M is a $d \times d$ integer matrix with determinant $(-1)^{d-1}$. It is easily checked that the matrix M is irreducible. Here a nonnegative matrix A is irreducible if for each ordered pair of indices I, J , there exists some $n \geq 0$ such that $A^n_{IJ} > 0$, where A_{IJ} means the (I, J) -element of the matrix A . An eigenvector α corresponding to the eigenvalue β of M and an eigenvector γ corresponding to β of the transpose of M are vectors $\alpha = {}^t[\alpha_1, \alpha_2, \dots, \alpha_d]$ and $\gamma = {}^t[\gamma_1, \gamma_2, \dots, \gamma_d]$, satisfying

$$(2.2) \quad M\alpha = \beta\alpha \quad \text{and} \quad {}^tM\gamma = \beta\gamma, \quad \text{respectively,}$$

where t indicates the transpose. From the Perron-Frobenius theory, irreducibility implies both eigenvectors are positive. We normalize α and γ by putting $\gamma_1 = 1$ and choosing α_i ($1 \leq i \leq d$) to satisfy $\langle \alpha, \gamma \rangle = 1$, where $\langle \cdot, \cdot \rangle$ denotes the standard inner product. By using (2.2), we can see that α_i and γ_i ($1 \leq i \leq d$) are given by

$$(2.3) \quad \alpha_i = \beta^{1-i} / \sum_{n=0}^{d-1} \beta^{-n} T^n_{\beta} 1,$$

$$\begin{aligned}
 \gamma_1 &= 1 = .k_1 k_2 \dots k_{d-1} 1, \\
 \gamma_2 &= T_\beta 1 = .k_2 \dots k_{d-1} 1, \\
 &\vdots \\
 \gamma_{d-1} &= T_\beta^{d-2} 1 = .k_{d-1} 1, \\
 \gamma_d &= T_\beta^{d-1} 1 = .1 = \frac{1}{\beta}.
 \end{aligned}
 \tag{2.4}$$

By $\mathbb{Z}[\beta]$ will be denoted the set of polynomials in β with integral coefficients. Then both $\{\alpha_1, \dots, \alpha_d\}$ and $\{\gamma_1, \dots, \gamma_d\}$ generate $\mathbb{Z}[\beta]$ and both are bases of $\mathbb{Q}(\beta)$.

It follows from either (2.4) or Theorem (Parry) in Section 1 that a sequence $(b_i)_{i=1}^\infty \in D_\beta$ if and only if for all i

$$0 \leq b_i \leq k_1,
 \tag{2.5}$$

$$\begin{cases}
 b_i = k_1 & \implies b_{i+1} \leq k_2, \\
 b_i = k_1, b_{i+1} = k_2 & \implies b_{i+2} \leq k_3, \\
 \vdots & \vdots \\
 b_i = k_1, b_{i+1} = k_2, \dots, b_{i+d-3} = k_{d-2} & \implies b_{i+d-2} \leq k_{d-1}, \\
 b_i = k_1, b_{i+1} = k_2, \dots, b_{i+d-3} = k_{d-2}, b_{i+d-2} = k_{d-1} & \implies b_{i+d-1} = 0.
 \end{cases}
 \tag{2.6}$$

Thus D_β is represented by the labeled graph \mathcal{G} in Figure 2. In other words, the admissible sequence $(b_i)_{i=1}^\infty$ of β -expansions is an infinite label of the walk in the sofic shift $X_{\mathcal{G}}$. See [12] concerning a labeled graph and a sofic shift.

§3. Substitutions

Let σ be the substitution of the alphabet $\mathcal{A} = \{1, 2, \dots, d\}$ given by:

$$\begin{aligned}
 \sigma : \quad 1 &\longrightarrow \underbrace{1 \dots 1}_{k_1} 2 \\
 &\quad \quad \quad 2 &\longrightarrow \underbrace{1 \dots 1}_{k_2} 3 \\
 &\quad \quad \quad \dots &\quad \quad \quad \dots \\
 &\quad \quad \quad d-1 &\longrightarrow \underbrace{1 \dots 1}_{k_{d-1}} d
 \end{aligned}$$




Figure 2: Labeled graph \mathcal{G} .

$$d \longrightarrow 1.$$

The free monoid on \mathcal{A} , that is to say, the set of finite words on \mathcal{A} , is denoted by $\mathcal{A}^* = \bigcup_{n=0}^{\infty} \mathcal{A}^n$.

There is a natural homomorphism (abelianization) $f : \mathcal{A}^* \rightarrow \mathbb{Z}^d$ given by $f(i) = \mathbf{e}_i$ for any $i \in \mathcal{A}$ where $\{\mathbf{e}_1, \dots, \mathbf{e}_d\}$ is the canonical basis of \mathbb{R}^d . Then there exists a unique linear transformation ${}^0\sigma$ satisfying the following commutative diagram:

$$\begin{array}{ccc} \mathcal{A}^* & \xrightarrow{\sigma} & \mathcal{A}^* \\ f \downarrow & & \downarrow f \\ \mathbb{Z}^d & \xrightarrow{{}^0\sigma} & \mathbb{Z}^d. \end{array}$$

We know that ${}^0\sigma$ is given by the matrix M in Section 2 in our case.

Let \mathcal{P} be the contractive invariant plane of M , that is,

$$\mathcal{P} = \{\mathbf{x} \in \mathbb{R}^d \mid \langle \mathbf{x}, \boldsymbol{\gamma} \rangle = 0\}.$$




Figure 3: The figure for ${}^1\sigma^*$ for the Rauzy fractal ($k_1 = k_2 = 1$).

Let $\pi : \mathbb{R}^d \rightarrow \mathcal{P}$ be the projection along the eigenvector α . In [15], Rauzy constructed a curious compact domain with a fractal boundary, called the Rauzy fractal, by using the Pisot number β for which $\text{Irr}(\beta) = x^3 - x^2 - x - 1$ ($k_1 = k_2 = 1$). Arnoux and Ito in [2] showed that for any Pisot substitution σ a compact domain X with a fractal boundary can be similarly constructed, using the following mapping ${}^1\sigma^*$:

$$(3.1) \quad {}^1\sigma^*(\mathbf{x}, i^*) = \sum_{j=1}^d \sum_{W_n^{(j)}=i} (M^{-1}(\mathbf{x} - f(P_n^{(j)})), j^*),$$

where $\sigma(j) = W_1^{(j)} \dots W_l^{(j)}$, $W_n^{(j)} \in \{1, \dots, d\}$, $P_n^{(j)}$ is the prefix of the letter $W_n^{(j)}$, and (\mathbf{x}, i^*) is the set $\{\mathbf{x} + \mathbf{e}_i + \sum_{j \neq i} \lambda_j \mathbf{e}_j \mid \lambda_j \in [0, 1]\}$. (See Figure 3.) We remark that we use the notation ${}^1\sigma^*$ in stead of $E_1^*(\sigma)$ which was used in [2].

In [17], the authors define higher dimensional extensions ${}^k\sigma$ ($1 \leq k \leq d$) of σ , acting on formal sums of weighted k -dimensional faces of unit cubes with vertices in \mathbb{Z}^d , and their dual maps ${}^k\sigma^*$. Moreover, they proved that these maps commute with the natural boundary morphisms and establish some basic properties.

THEOREM. *The following limit sets exist in the sense of Hausdorff metric:*

$$\begin{aligned} X_i &:= \lim_{n \rightarrow \infty} M^n(\pi({}^1\sigma^{*n}(\mathbf{0}, i^*))) \\ &= \lim_{n \rightarrow \infty} M^n(\pi({}^1\sigma^{*n}(-\mathbf{e}_i, i^*))), \quad (1 \leq i \leq d) \\ X &= \bigcup_{i=1}^d X_i. \end{aligned}$$

X_i are bounded, closed, and disjoint, up to a set of measure 0.




Figure 4: The Rauzy fractal ($k_1 = k_2 = 1$).

Note that the origin of \mathbb{R}^{d-1} belongs to X . (See the details in [2].) See Figure 4 in case $k_1 = k_2 = 1$.

From the equation (3.1) and $M^{-1}\mathbf{e}_1 = \mathbf{e}_d$, we see that the mapping ${}^1\sigma^*(\mathbf{0}, i^*)$ ($1 \leq i \leq d$) in our case are given by

$$\begin{aligned}
 {}^1\sigma^* : (\mathbf{0}, 1^*) &\longmapsto \sum_{i_1=0}^{k_1-1} (-i_1\mathbf{e}_d, 1^*) + \sum_{i_2=0}^{k_2-1} (-i_2\mathbf{e}_d, 2^*) \\
 &\quad + \cdots + \sum_{i_{d-1}=0}^{k_{d-1}-1} (-i_{d-1}\mathbf{e}_d, d-1^*) + (\mathbf{0}, d^*), \\
 (\mathbf{0}, 2^*) &\longmapsto (-k_1\mathbf{e}_d, 1^*), \\
 (\mathbf{0}, 3^*) &\longmapsto (-k_2\mathbf{e}_d, 2^*), \\
 &\quad \vdots \\
 (\mathbf{0}, d^*) &\longmapsto (-k_{d-1}\mathbf{e}_d, d-1^*).
 \end{aligned}$$

Hence, $M^{-1}X_i$ ($1 \leq i \leq d$) are given by

$$\begin{aligned}
 M^{-1}X_1 &= \lim_{n \rightarrow \infty} M^{n-1}\pi\left({}^1\sigma^{*n-1}({}^1\sigma^*(\mathbf{0}, 1^*))\right) \\
 &= \lim_{n \rightarrow \infty} M^{n-1}\pi\left({}^1\sigma^{*n-1}\left(\sum_{i_1=0}^{k_1-1} (-i_1\mathbf{e}_d, 1^*) + \cdots \right. \right. \\
 &\quad \left. \left. + \sum_{i_{d-1}=0}^{k_{d-1}-1} (-i_{d-1}\mathbf{e}_d, d-1^*) + (\mathbf{0}, d^*)\right)\right)
 \end{aligned}$$

$$= \bigcup_{i_1=0}^{k_1-1} (X_1 - i_1\pi\mathbf{e}_d) \cup \dots \cup \bigcup_{i_{d-1}=0}^{k_{d-1}-1} (X_{d-1} - i_{d-1}\pi\mathbf{e}_d) \cup X_d,$$

$$\begin{aligned} M^{-1}X_2 &= \lim_{n \rightarrow \infty} M^{n-1}\pi \left({}^1\sigma^{*n-1} ({}^1\sigma^*(\mathbf{0}, 2^*)) \right) \\ &= \lim_{n \rightarrow \infty} M^{n-1}\pi \left({}^1\sigma^{*n-1} (-k_1\mathbf{e}_d, 1^*) \right) \\ &= X_1 - k_1\pi\mathbf{e}_d, \\ &\quad \vdots \\ M^{-1}X_d &= \lim_{n \rightarrow \infty} M^{n-1}\pi \left({}^1\sigma^{*n-1} ({}^1\sigma^*(\mathbf{0}, d^*)) \right) \\ &= \lim_{n \rightarrow \infty} M^{n-1}\pi \left({}^1\sigma^{*n-1} (-k_{d-1}\mathbf{e}_d, d - 1^*) \right) \\ &= X_{d-1} - k_{d-1}\pi\mathbf{e}_d. \end{aligned}$$

Then applying M , from the property $M\pi\mathbf{e}_d = \pi M\mathbf{e}_d = \pi\mathbf{e}_1$, we have

$$(3.2) \quad \begin{cases} X_1 = \bigcup_{i_1=0}^{k_1-1} (MX_1 - i_1\pi\mathbf{e}_1) \cdots \bigcup_{i_{d-1}=0}^{k_{d-1}-1} (MX_{d-1} - i_{d-1}\pi\mathbf{e}_1) \cup MX_d, \\ X_2 = MX_1 - k_1\pi\mathbf{e}_1, \\ \quad \vdots \\ X_d = MX_{d-1} - k_{d-1}\pi\mathbf{e}_1, \end{cases}$$

$$(3.3) \quad \begin{aligned} X &= \bigcup_{i=1}^d X_i \\ &= \bigcup_{i_1=0}^{k_1} (MX_1 - i_1\pi\mathbf{e}_1) \cdots \bigcup_{i_{d-1}=0}^{k_{d-1}} (MX_{d-1} - i_{d-1}\pi\mathbf{e}_1) \cup MX_d. \end{aligned}$$

Since X_i are disjoint up to a set of measure 0, the partition of X is constructed. By using the partition (3.3), the transformation T_β^* on X without boundaries can be defined as follows:

$$(3.4) \quad T_\beta^*\mathbf{x} = M^{-1}\mathbf{x} + b^*\pi\mathbf{e}_d \quad \text{if } \mathbf{x} \in MX_j - b^*\pi\mathbf{e}_1 \text{ for some } j \text{ and } b^*.$$

Then for $\mathbf{x} \in X$ satisfying the condition that $T_\beta^{*k}x$ are not on the boundaries of X^i for any k , there exists an infinite sequence $(b_k^*)_{k=1}^\infty$ such that

$$(3.5) \quad T_\beta^{*k-1}\mathbf{x} \in MX_{j(k)} - b_k^*\pi\mathbf{e}_1,$$

and \mathbf{x} is represented by

$$\mathbf{x} = - \sum_{k=1}^\infty b_k^*M^{k-1}\pi\mathbf{e}_1.$$

Note that $(j(k))_{k=1}^\infty$ is the orbit of the point \mathbf{x} , that is,

$$T_\beta^{*k}\mathbf{x} \in X_{j(k)}.$$

From the set equations (3.2) and (3.5) we can see that

$$\begin{aligned} T_\beta^*(X_1^\circ) &= X_1^\circ \cup X_2^\circ \cup \dots \cup X_d^\circ, \\ T_\beta^*(X_2^\circ) &= X_1^\circ, \\ T_\beta^*(X_3^\circ) &= X_2^\circ, \\ &\vdots \\ T_\beta^*(X_d^\circ) &= X_{d-1}^\circ, \end{aligned}$$

where for each i X_i° is given by

$$X_i^\circ = \left\{ \mathbf{x} \in X_i \mid \begin{array}{l} T_\beta^{*k}\mathbf{x} \text{ are not on the boundaries of } X_j \text{ for any } k \\ \text{and any } j \end{array} \right\}.$$

Hence, an infinite walk $(b_k^*)_{k=1}^\infty$ is obtained from the labeled graph \mathcal{G}^* , which is the dual graph of \mathcal{G} . Here, the dual graph G^* is the graph with the same vertices as G , but with each edge in G reversed in direction. We can deal with all points of X_i successfully. As a consequence, we know that the domains X_i s ($1 \leq i \leq d$) are given by

$$(3.6) \quad X_i = \left\{ - \sum_{k=1}^\infty b_k^*M^{k-1}\pi\mathbf{e}_1 \mid \begin{array}{l} (b_k^*)_{k=1}^\infty \text{ is an admissible walk starting} \\ \text{at } i \text{ in } \mathcal{G}^* \end{array} \right\}.$$

See details in [2] and [6].




Figure 5: The figure of $\widehat{X} = \bigcup_{i=1}^d \widehat{X}_i$ in case $k_1 = k_2 = 1$.

§4. The natural extension of the β -transformation T_β

Let for each i ($1 \leq i \leq d$) $\widehat{X}_i \subset \mathbb{R}^d$ be the following domain:

$$(4.1) \quad \widehat{X}_i = \{t\alpha + \mathbf{x} \mid 0 \leq t < \gamma_i \text{ and } \mathbf{x} \in X_i\}.$$

And we define \widehat{X} by

$$\widehat{X} = \bigcup_{i=1}^d \widehat{X}_i.$$

See Figure 5.

Let \widehat{T}_β be the transformation on \widehat{X} given by

$$(4.2) \quad \widehat{T}_\beta(\underline{t}\alpha + \mathbf{x}) = \frac{(\beta t - [\beta t])\alpha + M\mathbf{x} - [\beta t]\pi\mathbf{e}_1}{T_\beta t}.$$

Note that \widehat{T}_β is just the β -transformation T_β on the direction α . We know that for any $\mathbf{z} = t\alpha + \mathbf{x} \in \widehat{X}$,

$$\widehat{T}_\beta(\mathbf{z}) = M\mathbf{z} - [\beta t]\mathbf{e}_1.$$

\widehat{T}_β will be a toral automorphism associated with M on the fundamental domain \widehat{X} .

From the partition (3.3) and the property (2.4) we have the following result.

PROPOSITION 4.1. \widehat{T}_β is surjective and injective on \widehat{X} except on the boundary.

Proof. For any $\mathbf{y} \in \widehat{X}$ there exist $\mathbf{y}' \in X_i$ ($1 \leq i \leq d$) and t' ($0 \leq t' < \gamma_i \leq 1$) such that

$$\mathbf{y} = t'\boldsymbol{\alpha} + \mathbf{y}'.$$

From the partition (3.2), we have

$$\mathbf{y} = t'\boldsymbol{\alpha} + M\mathbf{x}' - k\pi\mathbf{e}_1 \quad \text{for some } \mathbf{x}' \in X_j \text{ and } 0 \leq k \leq k_1.$$

Let

$$\mathbf{x} = \left(\frac{k}{\beta} + \frac{t'}{\beta} \right) \boldsymbol{\alpha} + \mathbf{x}'.$$

Then

$$\widehat{T}_\beta \mathbf{x} = t'\boldsymbol{\alpha} + M\mathbf{x}' - k\pi\mathbf{e}_1 = \mathbf{y}.$$

If $i = 1$, $0 \leq t' < 1$ and $k = 0, 1, \dots, k_j - 1$. Here we set $k_d = 1$. Then

$$0 \leq \frac{k}{\beta} + \frac{t'}{\beta} < \frac{k_j - 1}{\beta} + \frac{1}{\beta} = \frac{k_j}{\beta} = .k_j \leq \gamma_j.$$

If $i = 2, \dots, d$, we know that $0 \leq t' < \gamma_i = .k_i \dots k_{d-1}1$, $j = i - 1$, and $k = k_{i-1}$. Hence

$$0 \leq \frac{k}{\beta} + \frac{t'}{\beta} < \frac{k_{i-1}}{\beta} + \frac{\gamma_i}{\beta} = .k_{i-1} \dots k_{d-1}1 = \gamma_{i-1} = \gamma_j.$$

Therefore for any i , we see that $\mathbf{x} \in \widehat{X}_j \subset \widehat{X}$. Hence \widehat{T}_β is surjective. And except for the boundary, i, t', j , and k are uniquely determined by \mathbf{y} . Therefore \widehat{T}_β is almost everywhere injective. □

Therefore \widehat{T}_β is the natural extension of the transformation T_β .

Recall that the domain X is on the plane \mathcal{P} , which is orthogonal to $\boldsymbol{\gamma}$.

We put

$$Q := \begin{bmatrix} \alpha_1^{(1)} & \cdots & \alpha_1^{(r_1)} & \Re\alpha_1^{(r_1+1)} & -\Im\alpha_1^{(r_1+1)} & \cdots & \Re\alpha_1^{(r_1+r_2)} & -\Im\alpha_1^{(r_1+r_2)} \\ \alpha_2^{(1)} & \cdots & \alpha_2^{(r_1)} & \Re\alpha_2^{(r_1+1)} & -\Im\alpha_2^{(r_1+1)} & \cdots & \Re\alpha_2^{(r_1+r_2)} & -\Im\alpha_2^{(r_1+r_2)} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots & \vdots \\ \alpha_d^{(1)} & \cdots & \alpha_d^{(r_1)} & \Re\alpha_d^{(r_1+1)} & -\Im\alpha_d^{(r_1+1)} & \cdots & \Re\alpha_d^{(r_1+r_2)} & -\Im\alpha_d^{(r_1+r_2)} \end{bmatrix} \\ =: [\boldsymbol{\alpha}, \boldsymbol{\alpha}_2, \dots, \boldsymbol{\alpha}_d],$$

where \Re indicates the real part and \Im indicates the imaginary part. The plane \mathcal{P} is spanned by $\alpha_2, \alpha_3, \dots$, and α_d , because α_i ($2 \leq i \leq d$) and γ intersect orthogonally and α_i s are linearly independent.

Let us define the domains \widehat{Y} and \widehat{Y}_i ($1 \leq i \leq d$) as follows:

$$(4.3) \quad \widehat{Y} := Q^{-1}(\widehat{X}) \quad \text{and} \quad \widehat{Y}_i := Q^{-1}(\widehat{X}_i).$$

We will make preparations for the explicit representation of \widehat{Y}_i .

Define a $d \times d$ matrix

$$P := \begin{bmatrix} \alpha_1^{(1)} & \cdots & \alpha_1^{(r_1)} & \alpha_1^{(r_1+1)} & \overline{\alpha_1^{(r_1+1)}} & \cdots & \alpha_1^{(r_1+r_2)} & \overline{\alpha_1^{(r_1+r_2)}} \\ \alpha_2^{(1)} & \cdots & \alpha_2^{(r_1)} & \alpha_2^{(r_1+1)} & \overline{\alpha_2^{(r_1+1)}} & \cdots & \alpha_2^{(r_1+r_2)} & \overline{\alpha_2^{(r_1+r_2)}} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots & \vdots \\ \alpha_d^{(1)} & \cdots & \alpha_d^{(r_1)} & \alpha_d^{(r_1+1)} & \overline{\alpha_d^{(r_1+1)}} & \cdots & \alpha_d^{(r_1+r_2)} & \overline{\alpha_d^{(r_1+r_2)}} \end{bmatrix}$$

$$=: [\alpha, \mathbf{u}_2, \dots, \mathbf{u}_d].$$

Let

$$U := I_{r_1} \oplus \begin{bmatrix} 1 & \\ -\sqrt{-1} & \sqrt{-1} \end{bmatrix} \oplus \begin{bmatrix} 1 & \\ -\sqrt{-1} & \sqrt{-1} \end{bmatrix} \oplus \cdots \oplus \begin{bmatrix} 1 & \\ -\sqrt{-1} & \sqrt{-1} \end{bmatrix},$$

where $A \oplus B$ is a matrix of a form:

$$\begin{bmatrix} A & \mathbf{0} \\ \mathbf{0} & B \end{bmatrix}$$

and I_{r_1} is the identity matrix of size r_1 . Then

$$QU = P.$$

From $I_d = P \cdot P^{-1}$, we have

$$\mathbf{e}_1 = (P^{-1})_{11}\alpha + (P^{-1})_{21}\mathbf{u}_2 + \cdots + (P^{-1})_{d1}\mathbf{u}_d.$$

Each \mathbf{u}_i ($2 \leq i \leq d$) is also orthogonal to γ . Therefore,

$$\langle \mathbf{e}_1, \gamma \rangle = \langle (P^{-1})_{11}\alpha, \gamma \rangle = (P^{-1})_{11}\langle \alpha, \gamma \rangle = (P^{-1})_{11}.$$

It follows that

$$(P^{-1})_{11} = 1$$

and

$$P^{-1}\mathbf{e}_1 = {}^t[1, 1, \dots, 1].$$

You can see the detailed proof in [11]. According to the relation $QU = P$

$$Q^{-1}\mathbf{e}_1 = {}^t[\underbrace{1, 1, \dots, 1}_{r_1}, \underbrace{2, 0, \dots, 2, 0}_{2r_2}].$$

Moreover, from $I_d = Q \cdot Q^{-1}$

$$\mathbf{e}_1 = \boldsymbol{\alpha} + \boldsymbol{\alpha}_2 + \dots + \boldsymbol{\alpha}_{r_1} + 2\boldsymbol{\alpha}_{r_1+1} + 2\boldsymbol{\alpha}_{r_1+3} + \dots + 2\boldsymbol{\alpha}_{d-1}.$$

Since π is the projection along $\boldsymbol{\alpha}$,

$$\pi\mathbf{e}_1 = \boldsymbol{\alpha}_2 + \dots + \boldsymbol{\alpha}_{r_1} + 2\boldsymbol{\alpha}_{r_1+1} + 2\boldsymbol{\alpha}_{r_1+3} + \dots + 2\boldsymbol{\alpha}_{d-1}.$$

Hence

$$(4.4) \quad Q^{-1}\pi\mathbf{e}_1 = {}^t[0, \underbrace{1, 1, \dots, 1}_{r_1}, \underbrace{2, 0, \dots, 2, 0}_{2r_2}].$$

LEMMA 4.2. *The following relation holds:*

$$MQ = Q \begin{bmatrix} \beta & & & & \\ & \beta^{(2)} & & & \\ & & \ddots & & \\ & & & \beta^{(r_1)} & \\ & & & & \oplus \left[\begin{array}{cc} \Re\beta^{(r_1+1)} & -\Im\beta^{(r_1+1)} \\ \Im\beta^{(r_1+1)} & \Re\beta^{(r_1+1)} \end{array} \right] \oplus \dots \oplus \left[\begin{array}{cc} \Re\beta^{(r_1+r_2)} & -\Im\beta^{(r_1+r_2)} \\ \Im\beta^{(r_1+r_2)} & \Re\beta^{(r_1+r_2)} \end{array} \right] \end{bmatrix}.$$

Proof. The relation (2.2) implies that

$$MP = PD,$$

where D is the diagonal matrix

$$D = \begin{bmatrix} \beta & & & & & & & & \\ & \beta^{(2)} & & & & & & & \\ & & \ddots & & & & & & \\ & & & \beta^{(r_1)} & & & & & \\ & & & & \beta^{(r_1+1)} & & & & \\ & & & & & \overline{\beta^{(r_1+1)}} & & & \\ & & & & & & \ddots & & \\ & & & & & & & \beta^{(r_1+r_2)} & \\ & & & & & & & & \overline{\beta^{(r_1+r_2)}} \end{bmatrix}.$$

Then, from the relation $P = QU$,

$$MQU = QUD.$$

Using

$$U^{-1} = I_{r_1} \oplus \frac{1}{2} \begin{bmatrix} 1 & \sqrt{-1} \\ 1 & -\sqrt{-1} \end{bmatrix} \oplus \cdots \oplus \frac{1}{2} \begin{bmatrix} 1 & \sqrt{-1} \\ 1 & -\sqrt{-1} \end{bmatrix},$$

we have

$$\begin{aligned} Q^{-1}MQ &= UDU^{-1} \\ &= \begin{bmatrix} \beta & & & \\ & \beta^{(2)} & & \\ & & \ddots & \\ & & & \beta^{(r_1)} \end{bmatrix} \\ &\quad \oplus \begin{bmatrix} \Re\beta^{(r_1+1)} & -\Im\beta^{(r_1+1)} \\ \Im\beta^{(r_1+1)} & \Re\beta^{(r_1+1)} \end{bmatrix} \oplus \cdots \oplus \begin{bmatrix} \Re\beta^{(r_1+r_2)} & -\Im\beta^{(r_1+r_2)} \\ \Im\beta^{(r_1+r_2)} & \Re\beta^{(r_1+r_2)} \end{bmatrix}. \end{aligned}$$

□

Hereafter we represent \widehat{Y}_i s as the domains in $\mathbb{R} \times \mathbb{R}^{d-1}$.

PROPOSITION 4.3. *The domains \widehat{Y}_i s are given by*

$$\widehat{Y}_i = \left\{ \left(t, -\sum_{k=1}^{\infty} b_k^* R^{k-1} \mathbf{v} \right) \mid 0 \leq t < \gamma_i \text{ and } (b_k^*)_{k=1}^{\infty} \text{ is an admissible walk starting at } i \text{ in } \mathcal{G}^* \right\},$$

where

$$\begin{aligned} R &= \begin{bmatrix} \beta^{(2)} & & & \\ & \ddots & & \\ & & & \beta^{(r_1)} \end{bmatrix} \\ &\quad \oplus \begin{bmatrix} \Re\beta^{(r_1+1)} & -\Im\beta^{(r_1+1)} \\ \Im\beta^{(r_1+1)} & \Re\beta^{(r_1+1)} \end{bmatrix} \oplus \cdots \oplus \begin{bmatrix} \Re\beta^{(r_1+r_2)} & -\Im\beta^{(r_1+r_2)} \\ \Im\beta^{(r_1+r_2)} & \Re\beta^{(r_1+r_2)} \end{bmatrix} \end{aligned}$$

and

$$\mathbf{v} = {}^t \left[\underbrace{1, \dots, 1}_{r_1-1}, \underbrace{2, 0, \dots, 2, 0}_{2r_2} \right].$$

Proof. The definitions of \widehat{Y}_i , \widehat{X}_i , and X_i , that is, (4.3), (4.1), and (3.6), show that

$$\begin{aligned} \widehat{Y}_i &= Q^{-1}(\widehat{X}_i) \\ &= Q^{-1}\left\{t\alpha - \sum_{k=1}^{\infty} b_k^* M^{k-1} \pi \mathbf{e}_1 \mid (*)\text{-condition}\right\} \\ &= \left\{tQ^{-1}\alpha - \sum_{k=1}^{\infty} b_k^* Q^{-1} M^{k-1} \pi \mathbf{e}_1 \mid (*)\text{-condition}\right\}. \end{aligned}$$

By Lemma 4.2

$$Q^{-1}M^{k-1} = (\beta^{k-1} \oplus R^{k-1}) Q^{-1}.$$

And using (4.4), we have

$$\begin{aligned} \widehat{Y}_i &= \left\{t\mathbf{e}_1 - \sum_{k=1}^{\infty} b_k^* (\beta^{k-1} \oplus R^{k-1}) Q^{-1} \pi \mathbf{e}_1 \mid (*)\text{-condition}\right\} \\ &= \left\{\left(t, -\sum_{k=1}^{\infty} b_k^* R^{k-1} \mathbf{v}\right) \mid (*)\text{-condition}\right\}. \end{aligned}$$

Here (*)-condition means that $0 \leq t < \gamma_i$ and $(b_k^*)_{k=1}^{\infty}$ is an admissible walk starting at i in \mathcal{G}^* . Therefore we arrive at the conclusion of the assertion. □

Naturally, we can define a transformation \widehat{S}_β on \widehat{Y} as follows:

$$(4.5) \quad \widehat{S}_\beta := Q^{-1} \circ \widehat{T}_\beta \circ Q.$$

Then \widehat{S}_β is also a natural extension of T_β .

PROPOSITION 4.4. *The transformation \widehat{S}_β on \widehat{Y} is given by*

$$\widehat{S}_\beta(t, \mathbf{x}) = (\beta t - [\beta t], R\mathbf{x} - [\beta t]\mathbf{v})$$

and surjective.

Proof. From (4.5) and (4.2), which are definitions of \widehat{S}_β and \widehat{T}_β ,

$$\begin{aligned} \widehat{S}_\beta(t, \mathbf{x}) &:= Q^{-1} \circ \widehat{T}_\beta \circ Q(t\mathbf{e}_1 + 0 \oplus \mathbf{x}) \\ &= Q^{-1} \circ \widehat{T}_\beta(t\boldsymbol{\alpha} + Q(0 \oplus \mathbf{x})) \\ &= Q^{-1}((\beta t - [\beta t])\boldsymbol{\alpha} + MQ(0 \oplus \mathbf{x}) - [\beta t]\pi\mathbf{e}_1) \\ &= (\beta t - [\beta t], R\mathbf{x} - [\beta t]\mathbf{v}). \end{aligned}$$

Surjectivity of \widehat{S}_β is obtained by Proposition 4.1. □

§5. The reduction theorem

In this section, we introduce reduced numbers and show our main theorem.

Let $\widetilde{Y} (\subset \mathbb{R} \times \mathbb{R}^{d-1})$ be the following product space:

$$\widetilde{Y} := [0, 1) \times \mathbb{R}^{d-1}.$$

Let \widetilde{S}_β be the transformation on \widetilde{Y} defined by

$$\widetilde{S}_\beta(x, \mathbf{x}) := (\beta x - [\beta x], R\mathbf{x} - [\beta x]\mathbf{v}), \quad x \in [0, 1).$$

Then the restriction of \widetilde{S}_β on $\widehat{Y} (\subset \widetilde{Y})$ is \widehat{S}_β .

Define a map $\rho : \mathbb{Q}(\beta) \rightarrow \mathbb{R} \times \mathbb{R}^{d-1}$ by

$$\rho(x) = \left(x, \begin{bmatrix} x^{(2)} \\ \vdots \\ x^{(r_1)} \\ 2\Re x^{(r_1+1)} \\ 2\Im x^{(r_1+1)} \\ \vdots \\ 2\Re x^{(r_1+r_2)} \\ 2\Im x^{(r_1+r_2)} \end{bmatrix} \right).$$

DEFINITION 5.1. A real number $x \in \mathbb{Q}(\beta) \cap [0, 1)$ is reduced if $\rho(x) \in \widehat{Y}$.

In order to prove the main theorem, we will need some important lemmas.

LEMMA 5.1. *Let $x \in \mathbb{Q}(\beta) \cap [0, 1]$. Then*

$$\widetilde{S}_\beta(\rho(x)) = \rho(T_\beta x).$$

Proof. From the definitions of \widetilde{S}_β , ρ , and T_β , we have

$$\begin{aligned} \widetilde{S}_\beta(\rho(x)) &= \widetilde{S}_\beta \left(x, \begin{bmatrix} x^{(2)} \\ \vdots \\ x^{(r_1)} \\ 2\Re x^{(r_1+1)} \\ 2\Im x^{(r_1+1)} \\ \vdots \\ 2\Re x^{(r_1+r_2)} \\ 2\Im x^{(r_1+r_2)} \end{bmatrix} \right) \\ &= \left(\beta x - [\beta x], \begin{bmatrix} \beta^{(2)} x^{(2)} \\ \vdots \\ \beta^{(r_1)} x^{(r_1)} \\ 2(\Re \beta^{(r_1+1)} \cdot \Re x^{(r_1+1)} - \Im \beta^{(r_1+1)} \cdot \Im x^{(r_1+1)}) \\ 2(\Im \beta^{(r_1+1)} \cdot \Re x^{(r_1+1)} + \Re \beta^{(r_1+1)} \cdot \Im x^{(r_1+1)}) \\ \vdots \\ 2(\Re \beta^{(r_1+r_2)} \cdot \Re x^{(r_1+r_2)} - \Im \beta^{(r_1+r_2)} \cdot \Im x^{(r_1+r_2)}) \\ 2(\Im \beta^{(r_1+r_2)} \cdot \Re x^{(r_1+r_2)} + \Re \beta^{(r_1+r_2)} \cdot \Im x^{(r_1+r_2)}) \end{bmatrix} \right) \\ &\quad - [\beta x] \begin{bmatrix} 1 \\ \vdots \\ 1 \\ 2 \\ 0 \\ \vdots \\ 2 \\ 0 \end{bmatrix} \end{aligned}$$

using the relations $\mathfrak{R}(xy) = \mathfrak{R}x\mathfrak{R}y - \mathfrak{S}x\mathfrak{S}y$ and $\mathfrak{S}(xy) = \mathfrak{R}x\mathfrak{S}y + \mathfrak{S}x\mathfrak{R}y$,

$$\begin{aligned}
 &= \left(\beta x - [\beta x], \begin{bmatrix} (\beta x)^{(2)} - [\beta x] \\ \vdots \\ (\beta x)^{(r_1)} - [\beta x] \\ 2\mathfrak{R}((\beta x)^{(r_1+1)} - [\beta x]) \\ 2\mathfrak{S}((\beta x)^{(r_1+1)} - [\beta x]) \\ \vdots \\ 2\mathfrak{R}((\beta x)^{(r_1+r_2)} - [\beta x]) \\ 2\mathfrak{S}((\beta x)^{(r_1+r_2)} - [\beta x]) \end{bmatrix} \right) \\
 &= \rho(\beta x - [\beta x]) \\
 &= \rho(T_\beta x).
 \end{aligned}$$

Therefore we arrive at the conclusion. □

LEMMA 5.2. *Let $x \in \mathbb{Q}(\beta) \cap [0, 1)$ be reduced. Then*

- (1) $T_\beta x$ is reduced,
- (2) there exists x^* such that x^* is reduced and $T_\beta x^* = x$.

Proof. Since $x \in \mathbb{Q}(\beta) \cap [0, 1)$ is reduced, $\rho(x) \in \widehat{Y}$.

(1) From Lemma 5.1,

$$\widehat{S}_\beta(\rho(x)) = \rho(T_\beta x) \in \widehat{Y}.$$

Hence $T_\beta x$ is reduced.

(2) From Proposition 4.4, \widehat{S}_β is surjective on \widehat{Y} . Thus there exist $(x^*, \mathbf{x}) \in \widehat{Y}$ such that

$$(5.1) \quad \widehat{S}_\beta(x^*, \mathbf{x}) = \rho(x).$$

Comparing first coordinates in both sides, we see that

$$T_\beta x^* = x.$$

To verify x^* is reduced, we will only show

$$(x^*, \mathbf{x}) = \rho(x^*).$$

Then $\rho(x^*) \in \widehat{Y}$ implies that x^* is reduced.

We put

$$\mathbf{x} = {}^t [x_2, \dots, x_{r_1}, x_{r_1+1}, \widetilde{x_{r_1+1}}, \dots, x_{r_1+r_2}, \widetilde{x_{r_1+r_2}}].$$

Then (5.1) shows that

$$\beta x^* - [\beta x^*] = x$$

and

$$R\mathbf{x} - [\beta x^*]\mathbf{v} = \begin{bmatrix} x^{(2)} \\ \vdots \\ x^{(r_1)} \\ 2\Re x^{(r_1+1)} \\ 2\Im x^{(r_1+1)} \\ \vdots \\ 2\Re x^{(r_1+r_2)} \\ 2\Im x^{(r_1+r_2)} \end{bmatrix}.$$

So that,

$$\begin{bmatrix} \beta^{(2)}x_2 - [\beta x^*] \\ \vdots \\ \beta^{(r_1)}x_{r_1} - [\beta x^*] \\ \Re\beta^{(r_1+1)} \cdot x_{r_1+1} - \Im\beta^{(r_1+1)} \cdot \widetilde{x_{r_1+1}} - 2[\beta x^*] \\ \Im\beta^{(r_1+1)} \cdot x_{r_1+1} + \Re\beta^{(r_1+1)} \cdot \widetilde{x_{r_1+1}} \\ \vdots \\ \Re\beta^{(r_1+r_2)} \cdot x_{r_1+r_2} - \Im\beta^{(r_1+r_2)} \cdot \widetilde{x_{r_1+r_2}} - 2[\beta x^*] \\ \Im\beta^{(r_1+r_2)} \cdot x_{r_1+r_2} + \Re\beta^{(r_1+r_2)} \cdot \widetilde{x_{r_1+r_2}} \end{bmatrix} = \begin{bmatrix} \beta^{(2)}x^{*(2)} - [\beta x^*] \\ \vdots \\ \beta^{(r_1)}x^{*(r_1)} - [\beta x^*] \\ 2\Re \left(\beta^{(r_1+1)}x^{*(r_1+1)} - [\beta x^*] \right) \\ 2\Im \left(\beta^{(r_1+1)}x^{*(r_1+1)} - [\beta x^*] \right) \\ \vdots \\ 2\Re \left(\beta^{(r_1+r_2)}x^{*(r_1+r_2)} - [\beta x^*] \right) \\ 2\Im \left(\beta^{(r_1+r_2)}x^{*(r_1+r_2)} - [\beta x^*] \right) \end{bmatrix}.$$

Thus

$$x_2 = x^{*(2)}, \dots, x_{r_1} = x^{*(r_1)},$$

and for $1 \leq j \leq r_2$

$$\begin{aligned} & \Re\beta^{(r_1+j)} \cdot x_{r_1+j} - \Im\beta^{(r_1+j)} \cdot \widetilde{x_{r_1+j}} \\ &= 2\Re\left(\beta^{(r_1+j)}x^{*(r_1+j)}\right) \\ &= 2\Re\beta^{(r_1+j)}\Re x^{*(r_1+j)} - 2\Im\beta^{(r_1+j)}\Im x^{*(r_1+j)}, \\ & \Im\beta^{(r_1+j)} \cdot x_{r_1+j} + \Re\beta^{(r_1+j)} \cdot \widetilde{x_{r_1+j}} \\ &= 2\Im\left(\beta^{(r_1+j)}x^{*(r_1+j)}\right) \\ &= 2\Re\beta^{(r_1+j)}\Im x^{*(r_1+j)} + 2\Im\beta^{(r_1+j)}\Re x^{*(r_1+j)}. \end{aligned}$$

Then for $1 \leq j \leq r_2$, we have

$$\begin{aligned} & \left(x_{r_1+j} - 2\Re x^{*(r_1+j)}\right)\Re\beta^{(r_1+j)} - \left(\widetilde{x_{r_1+j}} - 2\Im x^{*(r_1+j)}\right)\Im\beta^{(r_1+j)} = 0, \\ & \left(x_{r_1+j} - 2\Re x^{*(r_1+j)}\right)\Im\beta^{(r_1+j)} + \left(\widetilde{x_{r_1+j}} - 2\Im x^{*(r_1+j)}\right)\Re\beta^{(r_1+j)} = 0. \end{aligned}$$

Thus

$$x_{r_1+j} = 2\Re x^{*(r_1+j)} \quad \text{and} \quad \widetilde{x_{r_1+j}} = 2\Im x^{*(r_1+j)}.$$

Therefore

$$(x^*, \mathbf{x}) = \rho(x^*).$$

Thus we obtain the assertion (2). □

By the lemmas above, we can get a sufficient condition for pure periodicity of β -expansions.

PROPOSITION 5.3. *Let $x \in \mathbb{Q}(\beta) \cap [0, 1)$ be reduced. Then x has a purely periodic β -expansion.*

Proof. Lemma 5.2 (2) shows that there exist x_i^* 's such that x_i^* 's are reduced and $T_\beta x_i^* = x_{i-1}^*$, where we set $x_0^* = x$. Here, we put

$$x = \frac{p_0}{q} \quad \text{for some } q \in \mathbb{Z}, p_0 \in \mathbb{Z}[\beta].$$

Then $T_\beta x_1^* = x$ implies that

$$\beta x_1^* - [\beta x_1^*] = x.$$

So that

$$x_1^* = \frac{[\beta x_1^*]}{\beta} + \frac{x}{\beta} = \frac{p_1}{q} \text{ for some } p_1 \in \mathbb{Z}[\beta].$$

Inductively we can see for every k

$$x_k^* = \frac{p_k}{q} \text{ for some } p_k \in \mathbb{Z}[\beta].$$

Let b_j be positive real numbers. Only in this proof, we denote by $x^{(j)}$ ($1 \leq j \leq d$) algebraic conjugates of x and $x^{(1)} = x$. Let

$$C = \{x \in \mathbb{Z}[\beta] \mid |x^{(j)}| \leq b_j\}.$$

Obviously, C is a finite set. As \widehat{Y} is bounded, we can see the set $\{x_i^*\}_{i=0}^\infty$ is a finite set. Hence there exist j and k ($j > k$) such that

$$x_j^* = x_{j-k}^*.$$

Applying T_β^{j-k} we get

$$x_k^* = x.$$

Hence

$$T_\beta^k x = x.$$

Therefore x has a purely periodic β -expansion. □

Lemma 5.2 and Proposition 5.3 show that the transformation T_β restricted to $\mathbb{Q}(\beta) \cap [0, 1)$ is bijective.

To complete the proof of our main theorem, the following proposition is positively necessary.

PROPOSITION 5.4. *Let $x \in \mathbb{Q}(\beta) \cap [0, 1)$. Then there exists $N_1 > 0$ such that $T_\beta^N x$ are reduced for any $N \geq N_1$.*

Proof. Consider the Euclidean distance d between $\widetilde{S}_\beta^k(\rho(x))$ and $\widetilde{S}_\beta^k(x, \mathbf{0})$ for $k \geq 0$. Since the first coordinates of these points are equal,

this coincides with the distance between the origin of \mathbb{R}^{d-1} and

$$R^k \begin{bmatrix} x^{(2)} \\ \vdots \\ x^{(r_1)} \\ 2\Re x^{(r_1+1)} \\ 2\Im x^{(r_1+1)} \\ \vdots \\ 2\Re x^{(r_1+r_2)} \\ 2\Im x^{(r_1+r_2)} \end{bmatrix}.$$

By $s(x)$ we denote this distance. As R^k s are given by

$$R^k = \begin{bmatrix} (\beta^{(2)})^k & & \\ & \ddots & \\ & & (\beta^{(r_1)})^k \end{bmatrix} \\ \oplus |\beta^{(r_1+1)}|^{2k} \begin{bmatrix} \Re \beta^{(r_1+1)} / |\beta^{(r_1+1)}|^2 & -\Im \beta^{(r_1+1)} / |\beta^{(r_1+1)}|^2 \\ \Im \beta^{(r_1+1)} / |\beta^{(r_1+1)}|^2 & \Re \beta^{(r_1+1)} / |\beta^{(r_1+1)}|^2 \end{bmatrix}^k \\ \oplus \dots \oplus |\beta^{(r_1+r_2)}|^{2k} \begin{bmatrix} \Re \beta^{(r_1+r_2)} / |\beta^{(r_1+r_2)}|^2 & -\Im \beta^{(r_1+r_2)} / |\beta^{(r_1+r_2)}|^2 \\ \Im \beta^{(r_1+r_2)} / |\beta^{(r_1+r_2)}|^2 & \Re \beta^{(r_1+r_2)} / |\beta^{(r_1+r_2)}|^2 \end{bmatrix}^k,$$

we have

$$s(x)^2 = (\beta^{(2)})^{2k} (x^{(2)})^2 + \dots + (\beta^{(r_1)})^{2k} (x^{(r_1)})^2 \\ + |\beta^{(r_1+1)}|^{2k} \left\{ (2\Re x^{(r_1+1)})^2 + (2\Im x^{(r_1+1)})^2 \right\} \\ + \dots + |\beta^{(r_1+r_2)}|^{2k} \left\{ (2\Re x^{(r_1+r_2)})^2 + (2\Im x^{(r_1+r_2)})^2 \right\}.$$

If we put

$$u = \max \{ |\beta^{(2)}|, \dots, |\beta^{(r_1)}|, |\beta^{(r_1+1)}|, \dots, |\beta^{(r_1+r_2)}| \},$$

then $0 < u < 1$ and

$$s(x) \leq u^k \cdot \left\{ (x^{(2)})^2 + \dots + (x^{(r_1)})^2 + (2\Re x^{(r_1+1)})^2 + (2\Im x^{(r_1+1)})^2 \right. \\ \left. + \dots + (2\Re x^{(r_1+r_2)})^2 + (2\Im x^{(r_1+r_2)})^2 \right\}^{1/2}.$$

Thus

$$d(\widetilde{S}_\beta^k(\rho(x)), \widetilde{S}_\beta^k(x, \mathbf{0})) \leq u^k \cdot d(\rho(x), (x, \mathbf{0})).$$

From the fact $(x, \mathbf{0}) \in \widehat{Y}$ and $\widetilde{S}_\beta|_{\widehat{Y}} = \widehat{S}_\beta$, we know that

$$\widetilde{S}_\beta^k(x, \mathbf{0}) \in \widehat{Y}.$$

It follows that $\widetilde{S}_\beta^k(\rho(x))$ comes exponentially close to \widehat{Y} as $k \rightarrow \infty$. By the same reason that we used in the proof of Proposition 5.3, we can conclude that there exists a finite number of $\rho(T_\beta^k)$ in a certain bounded domain. Hence

$$\widetilde{S}_\beta^{N_1}(\rho(x)) = \rho(T_\beta^{N_1}x) \in \widehat{Y}$$

for a sufficiently large N_1 . Then $T_\beta^{N_1}x$ is reduced. From Lemma 5.2 (1), we see that $T_\beta^N x$ are reduced for any $N \geq N_1$.

At last we attain our goal.

THEOREM 5.5. *Let $x \in [0, 1)$. Then*

- (1) $x \in \mathbb{Q}(\beta)$ if and only if x has an eventually periodic β -expansion,
- (2) $x \in \mathbb{Q}(\beta)$ is reduced if and only if x has a purely periodic β -expansion.

Proof. (1) Assume that $x \in \mathbb{Q}(\beta)$. By Proposition 5.4, there exists $N > 0$ such that $T_\beta^N x$ is reduced. Proposition 5.3 says that $T_\beta^N x$ has a purely periodic β -expansion. Hence x has an eventually periodic β -expansion. The opposite implication is trivial.

(2) Necessity is obtained by Proposition 5.3. Conversely, assume that x has a purely periodic β -expansion. According to Proposition 5.4, there exists $N > 0$ such that $T_\beta^N x$ is reduced. The pure periodicity of x implies that there exists $j > 0$ such that $T_\beta^{N+j} x = x$. Lemma 5.2 (1) says that x is reduced. \square

REFERENCES

- [1] S. Akiyama, Pisot numbers and greedy algorithm, Number Theory, Diophantine, Computational and Algebraic Aspects, (K. Györy, A. Pethö and V. T. Sós, eds.), de Gruyter, 1998, pp. 9–21.

- [2] P. Arnoux and Sh. Ito, *Pisot substitutions and Rauzy fractals*, **8** (2001), Bull. Belg. Math. Soc., 181–207.
- [3] A. Bertrand, *Développements en base de Pisot et répartition modulo 1*, **285** (1977), C.R. Acad. Sci, Paris, 419–421.
- [4] A. Brauer, *On algebraic equations with all but one root in the interior of the unit circle*, Math. Nachr., **4** (1951), 250–257.
- [5] F. M. Dekking, *Recurrent sets*, Adv. in Math., **44** (1982), 78–104.
- [6] H. Ei and Sh. Ito, *Tilings from characteristic polynomials of β -expansions*, preprint.
- [7] C. Frougny and B. Solomyak, *Finite beta-expansions*, Ergod. Th. and Dynam. Sys., **12** (1992), 713–723.
- [8] Y. Hara and Sh. Ito, *On real quadratic fields and periodic expansions*, Tokyo J. Math., **12** (1989), 357–370.
- [9] Sh. Ito, *On periodic expansions of cubic numbers and Rauzy fractals*, preprint.
- [10] Sh. Ito and M. Ohtsuki, *Modified Jacobi-Pirron algorithm and generating Markov partitions for special hyperbolic toral automorphisms*, Tokyo J. Math., **16** (1993), 441–472.
- [11] Sh. Ito and Y. Sano, *On periodic β -expansions of Pisot numbers and Rauzy fractals*, Osaka J. Math., **38** (2001), 349–368.
- [12] D. Lind and B. Marcus, *An Introduction to Symbolic Dynamics and Coding*, Cambridge Univ. Press, Cambridge, 1995.
- [13] W. Parry, *On the β -expansions of real numbers*, Acta Math. Acad. Sci. Hungar., **11** (1960), 401–416.
- [14] B. Praggastis, *Numeration systems and Markov partitions from self similar tilings*, Transactions of the American Mathematical Society, **351** (1999), 3315–3349.
- [15] G. Rauzy, *Nombres algébriques et substitutions*, Bull. Soc. Math. France, **110** (1982), 147–178.
- [16] A. Rényi, *Representations for real numbers and their ergodic properties*, Acta Math. Acad. Sci. Hungar., **8** (1957), 477–493.
- [17] Y. Sano, P. Arnoux, and Sh. Ito, *Higher dimensional extensions of substitutions and their dual maps*, J. d'Analyse Math., **83** (2001), 183–206.
- [18] K. Schmidt, *On periodic expansions of Pisot numbers and Salem numbers*, Bull. London math. Soc., **12** (1980), 269–278.
- [19] B. Solmyak, *Dynamics of self-similar tilings*, Ergod. Th. & Dynam. Sys., **17** (1997), 1–44.

Department of Mathematics and Computer Science
Tsuda College
2-1-1 Tsuda-Machi
Kodaira
Tokyo, 187-8577
Japan
`sano@tsuda.ac.jp`