

GENERALISED DISCRIMINANTS

C.F. WOODCOCK AND P.R. GRAVES-MORRIS

In this paper, we establish the generalised discriminant theorem governing uniqueness of symmetric polynomials in n variables of least total degree $d_{n,k}$ which vanish when any $k + 1$ of the variables are equal.

1. INTRODUCTION

The symmetric polynomial in the variables x_1, x_2, \dots, x_n of least total degree which vanishes when any two of these variables are equal is

$$P_{n,1}(x) = \prod_{1 \leq i < j \leq n} (x_i - x_j)^2,$$

where $x := (x_1, x_2, \dots, x_n)$. This polynomial is called the discriminant polynomial [2]. It has total degree $n(n - 1)$ and degree $2(n - 1)$ in each of the x_i .

Equation (20) gives the explicit formula which generalises $P_{n,1}(x)$ and is the (non-trivial) symmetric polynomial of least total degree in x_1, x_2, \dots, x_n which vanishes when any $k + 1$ of the x_i are equal. We call this polynomial the generalised discriminant and we show that it is unique up to an overall scale factor. These new results are summarised by Theorem 1.

2. GENERALISED DISCRIMINANTS

We shall show that for each $n, k \geq 1$ there is, up to a constant multiple, a unique (non-zero) symmetric polynomial $P_{n,k}(x_1, x_2, \dots, x_n)$ of least total degree (with coefficients in a field of characteristic zero) which vanishes whenever $k + 1$ of the x_i are equal. We shall further show that

- (i) $P_{n,k}(x_1, x_2, \dots, x_n)$ is a homogeneous polynomial,
- (ii) $\deg\{P_{n,k}\} = kq(q - 1) + 2rq$, where

$$(1) \quad q = [n/k] \text{ and } r = n - qk,$$

Received 20 April 1995.

P.R. Graves-Morris gratefully acknowledges an Erskine fellowship at Canterbury University. We are grateful to Dr. J.R. Merriman and Mr. R. Thukral for some helpful discussions and symbolic calculations.

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/96 \$A2.00+0.00.

and

(iii) $P_{n,k}(x_1, x_2, \dots, x_n)$ has degree $2[(n - 1)/k]$ in each variable x_i .

For brevity, and where appropriate, we let $\mathbf{x} := (x_1, x_2, \dots, x_n)$ denote the variables.

EXAMPLE 1. Case $k = 1$. In the introduction, we observed that

$$(2) \quad P_{n,1}(\mathbf{x}) = \prod_{1 \leq i < j \leq n} (x_i - x_j)^2$$

is called the discriminant polynomial in n variables. It is the symmetric polynomial of least degree which vanishes when any two of $\{x_i\}$ are equal and it is totally symmetric with respect to interchange of $\{x_i\}$. It is unique up to a non-zero constant multiplier. We note that $P_{n,1}(\mathbf{x})$ is of total degree $n(n - 1)$ in accord with (ii) and it is of degree $2(n - 1)$ in each x_i individually, in accord with (iii) above.

Before going on to the next example, we show that if $P(\mathbf{x})$ is to be a generalised discriminant polynomial of degree d say, then it is necessarily homogeneous. Consider the effect of a dilation $\mathbf{x} \rightarrow \alpha\mathbf{x}$ under which

$$P(\mathbf{x}) \rightarrow P(\mathbf{x}; \alpha) \equiv \sum_{j=0}^d p_j(\mathbf{x})\alpha^j.$$

$P(\mathbf{x}; \alpha)$ is a polynomial in the variables \mathbf{x}, α and it vanishes whenever $k + 1$ of its variables x_i are equal; each $p_j(\mathbf{x})$ inherits this property. By construction, each $p_j(\mathbf{x})$ is of total degree j . But, by hypothesis, d is minimal, and so $p_j(\mathbf{x}) \equiv 0$ for $j < d$. Consequently $P(\mathbf{x})$ and each $P_{n,k}(\mathbf{x})$ must be homogeneous polynomials.

EXAMPLE 2. Case $n = 3, k = 2$.

$$P_{3,2}(\mathbf{x}) = (x_1 - x_2)^2 + (x_2 - x_3)^2 + (x_3 - x_1)^2$$

is a non-trivial symmetric polynomial in x_1, x_2, x_3 and it vanishes identically if $x_1 = x_2 = x_3$. The only polynomial of lesser degree with these properties is $P(\mathbf{x}) = 0$. Because the required polynomial has to be homogeneous, it readily follows that $P_{3,2}(\mathbf{x})$ is unique up to an overall constant multiplier.

CONSTRUCTION OF THE GENERALISED DISCRIMINANT $P_{n,k}(\mathbf{x})$: Let the index set $\{1, 2, \dots, n\}$ denote the vertices of a graph G , and let G be the disjoint union of cliques C_1, C_2, \dots, C_k , where C_1, C_2, \dots, C_r each have $q + 1$ vertices and $C_{r+1}, C_{r+2}, \dots, C_k$ each have q vertices. (A clique with q vertices is the complete subgraph formed with these vertices [6].) Let E denote the set of all ordered pairs

(i, j) in which $i < j$ and i, j belong to the same clique. Thus E is the set of (directed) edges of G , and we form

$$(3) \quad P_E(\mathbf{x}) := \prod_{\ell=1}^k \prod_{\substack{i < j \\ i, j \in C_\ell}} (x_i - x_j)^2 = \prod_{(i, j) \in E} (x_i - x_j)^2.$$

We construct the generalised discriminant $P_{n,k}(\mathbf{x})$ as the symmetrised form of $p_E(\mathbf{x})$,

$$(4) \quad P_{n,k}(\mathbf{x}) := \sum_{\sigma \in S_n} \prod_{(i, j) \in E} (x_{\sigma(i)} - x_{\sigma(j)})^2$$

where S_n is the group of all permutations of $\{1, 2, \dots, n\}$.

It is easily verified that $P_{n,k}(\mathbf{x})$ defined by (4) possesses the properties (i), (ii), (iii) above. Property (i) is obvious. Property (ii) follows by counting the contributions $q(q + 1)$ to the degree from each clique C_1, \dots, C_r and $q(q - 1)$ from each clique C_{r+1}, \dots, C_k . We denote the degree of $P_{n,k}(\mathbf{x})$ by

$$(5) \quad \lambda_{n,k} := kq(q - 1) + 2rq.$$

Property (iii) follows by checking the power of x_1 and considering separately the two cases of $r = 0$ and $r > 0$.

It is also true that (4) expresses $P_{n,k}(\mathbf{x})$ as a sum of squares, each of which vanishes if any $k + 1$ of the x_i are equal. To see that $P_{n,k}(\mathbf{x})$ does not vanish identically when k or fewer of the x_i are equal, re-order the x_i so that variables equal by value occur consecutively. Place the variables one by one in this order into the cliques C_1, C_2, \dots, C_k consecutively. In this way, the cliques have their prescribed numbers of elements, no clique contains two variables equal by value, and a non-zero term of the form (3) has been constructed.

Next we must prove that $P_{n,k}(\mathbf{x})$ is of minimal degree. To this end, we use the characterisation of $P_{n,k}(\mathbf{x})$ given in Lemma 1, and we need some preliminary definitions.

DEFINITION: We define a *lexicographical order* \geq on the set of monomials $x_1^{s_1} x_2^{s_2} \dots x_n^{s_n}$ (each $s_i \geq 0$) by defining $x_1^{\ell_1} x_2^{\ell_2} \dots x_n^{\ell_n} \geq x_1^{m_1} x_2^{m_2} \dots x_n^{m_n}$ with equality if $\ell_i = m_i$ $i = 1, 2, \dots, n$ or inequality if $\ell_i > m_i$, where i is the least index with $\ell_i \neq m_i$.

DEFINITION: We define the *leading monomial* $\mathcal{M}(P)$ of a given polynomial $P(\mathbf{x})$ to be the greatest monomial (under the lexicographical order) with non-zero coefficient in $P(\mathbf{x})$.

Thus $\mathcal{M}(P)$ takes the form

$$(6) \quad \mathcal{M}(P) = x_1^{\ell_1} x_2^{\ell_2} \dots x_n^{\ell_n} \text{ with } \ell_1 \geq \ell_2 \geq \dots \geq \ell_n$$

if $P(\mathbf{x})$ is symmetric in x_1, x_2, \dots, x_n . The integers $(\ell_1, \ell_2, \dots, \ell_n)$ provide a partial characterisation of $P(\mathbf{x})$, and $P(\mathbf{x})$ obviously has degree ℓ_1 in the variable x_1 alone and, for that matter, degree ℓ_1 in each x_i if $P(\mathbf{x})$ is symmetric. We also have a *hierarchical property*: if

$$(7) \quad P(\mathbf{x}) = x_1^{\ell_1} Q(x_2, \dots, x_n) + \text{lower order terms in } x_1,$$

then

$$(8) \quad \mathcal{M}(P) = x_1^{\ell_1} \mathcal{M}(Q).$$

DEFINITION OF THE CHARACTER $H_{n,k}(\ell_1, \ell_2, \dots, \ell_n)$: If a polynomial $P(\mathbf{x})$ has the properties that

- (a) $P(\mathbf{x})$ is homogeneous,
- (b) $P(\mathbf{x})$ is symmetric in x_1, x_2, \dots, x_n ,
- (c) $P(\mathbf{x}) = 0$ whenever $k + 1$ of the x_i are equal,

then $P(\mathbf{x})$ is said to have character $H_{n,k}(\ell_1, \ell_2, \dots, \ell_n)$ where $\ell_1, \ell_2, \dots, \ell_n$ are the powers in the leading monomial $x_1^{\ell_1} \dots x_n^{\ell_n}$ of $P(\mathbf{x})$.

LEMMA 1. The generalised discriminant $P_{n,k}(\mathbf{x})$ defined by (4) has character $H_{n,k}(\ell_1, \ell_2, \dots, \ell_n)$ where

$$(9) \quad \ell_i := \lambda_{n-i+1,k} - \lambda_{n-i,k} = 2 \left[\frac{n-i}{k} \right]$$

and

$$(10) \quad \ell_1 + \ell_2 + \dots + \ell_n = \lambda_{n,k}.$$

PROOF: The remarks associated with the Definition (5) were made to establish that $\ell_1 = \lambda_{n,k} - \lambda_{n-1,k} = 2[(n-1)/k]$. Equation (9) follows from (5) together with the hierarchical principle expressed by (7), and (10) reflects the decomposition (6). \square

In order to prove that $P_{n,k}(\mathbf{x})$ is of minimal degree, we consider an arbitrary polynomial $P(\mathbf{x})$ which satisfies condition (a), (b) and (c) above. Then $P(\mathbf{x})$ has total degree $\ell_1 + \ell_2 + \dots + \ell_n$, it is of degree ℓ_1 in each x_i separately, and so $P(\mathbf{x})$ has character $H_{n,k}(\ell_1, \dots, \ell_n)$. If we express

$$(11) \quad P(\mathbf{x}) = x_1^{\ell_1} Q(x_2, \dots, x_n) + \text{lower order terms in } x_1,$$

then $Q(\mathbf{x})$ has character $H_{n-1,k}(\ell_2, \dots, \ell_n)$ in the variables x_2, \dots, x_n . This decomposition follows directly from the preceding definition of the character $H_{n,k}(\ell_1, \dots, \ell_n)$ and from its hierarchical property.

Similarly, we may express

$$(12) \quad P(\mathbf{x}) = x_1^{\ell_1} x_2^{\ell_2} \dots x_k^{\ell_k} R(x_{k+1}, \dots, x_n) + \text{lower order terms in } x_1, x_2, \dots, x_k$$

where $R(x_{k+1}, \dots, x_n)$ has character $H_{n-k,k}(\ell_{k+1}, \dots, \ell_n)$.

EXAMPLE 3.

$$P_{3,1}(x) = (x_1 - x_2)^2(x_1 - x_3)^2(x_2 - x_3)^2$$

is *Example 1* in the case $n = 3$. Here we have $\mathcal{M}(P_{3,1}) = x_1^4 x_2^2 x_3^0$ and $\ell_1 = 4, \ell_2 = 2, \ell_3 = 0$. We may also express

(13)

$$\begin{aligned} P_{3,1}(x) &= \sum_{m_1=0}^4 \sum_{m_2=0}^4 x_1^{m_1} x_2^{m_2} P_{m_1 m_2}(x_3) \\ &= x_1^4 x_2^2 - 2x_1^3 x_2^3 + x_1^2 x_2^4 + \text{terms of lower total degree in } x_1, x_2. \end{aligned}$$

We next establish several further properties of an arbitrary polynomial $P(x)$ satisfying conditions (a), (b), (c) above.

LEMMA 2. Suppose that $P(x)$ has character $H_{n,k}(\ell_1, \dots, \ell_n)$, and that $n \geq k + 1$. Then $\ell_1 \geq \ell_{k+1} + 2$.

PROOF: As in the previous example, express $P(x)$ as

$$(14) \quad P(x) = \sum_{\text{each } m_i \geq 0} P_{m_1, m_2, \dots, m_{k+1}}(x_{k+2}, \dots, x_n) x_1^{m_1} x_2^{m_2} \dots x_{k+1}^{m_{k+1}}$$

thereby defining the coefficients $P_{m_1, \dots, m_{k+1}}(x_{k+2}, \dots, x_n)$.

Since $P(x)$ is totally symmetric, each $P_{m_1, \dots, m_{k+1}}(x_{k+2}, \dots, x_n)$ depends only on the values $\{m_1, m_2, \dots, m_{k+1}\}$ and not on the order of these integers. From the hierarchical principle, the leading term of $P(x_1, x_2, \dots, x_n)$ is

$$x_1^{\ell_1} x_2^{\ell_2} \dots x_{k+1}^{\ell_{k+1}} P_{\ell_1, \dots, \ell_{k+1}}(x_{k+2}, \dots, x_n).$$

The characteristic property (c) above implies that

$$P(x, x, \dots, x, x_{k+2}, \dots, x_n) \equiv 0$$

and hence its leading coefficient in the variable x is zero. It is essential that this leading term has at least two source terms of the type displayed in (13). Otherwise, the characteristic property (c) above would imply that the leading term would vanish, and so would $P(x)$. We use this argument below.

Now suppose that the lemma is false. Then either

- (i) $\ell_1 = \ell_2 = \dots = \ell_{k+1} = \ell$ (say), or
- (ii) $\ell_1 = \ell_2 = \dots = \ell_s = \ell$ (say), and

$$\ell_{s+1} = \ell_{s+2} = \dots = \ell_{k+1} = \ell - 1 \text{ for some } s \in [1, k].$$

CASE (I). There is precisely one source term

$$P_{\ell, \ell, \dots, \ell}(x_{k+2}, \dots, x_n) x_1^\ell x_2^\ell \dots x_{k+1}^\ell$$

in (14) contributing to the leading monomial $x^{(k+1)\ell}$. The characteristic property (c) above implies that this term is zero, contradicting the hypothesis that it is a leading term. Therefore Case (i) cannot hold.

CASE (II). There are now several source terms in (14) contributing to the coefficient of $x^{\ell_1 + \ell_2 + \dots + \ell_{k+1}}$ in $P(x, \dots, x, x_{k+2}, \dots, x_n)$, which is clearly the leading monomial in x , and in fact they all provide equal contributions. The leading monomial of $P(x)$ is

$$(15) \quad t(x) := x_1^\ell x_2^\ell \dots x_s^\ell x_{s+1}^{\ell-1} \dots x_{k+1}^{\ell-1} P_{\underbrace{\ell, \ell, \dots, \ell}_s \text{ indices}, \underbrace{\ell-1, \dots, \ell-1}_{k+1-s} \text{ indices}}(x_{k+2}, \dots, x_n).$$

By permuting x_1, \dots, x_{k+1} in $t(x)$, and because the P coefficient in the right-hand side of (15) is symmetrical with respect to the order of its subscripts, we only find terms which make a contribution to $P(x, x, \dots, x, x_{k+2}, \dots, x_n)$ of $x^{\ell_1 + \ell_2 + \dots + \ell_{k+1}}$ $P_{\ell_1, \dots, \ell_1, \ell-1, \dots, \ell-1}(x_{k+2}, \dots, x_n)$.

The characteristic property (c) above implies that the total contribution is zero, contradicting the hypothesis that $t(x)$ in (15) is a leading term and so is non-zero. Therefore Case (ii) cannot hold. □

LEMMA 3. *Suppose that $P(x)$ has character $H_{n,k}(\ell_1, \dots, \ell_n)$. Then*

$$(16) \quad \ell_1 \geq 2 \left\lfloor \frac{n-1}{k} \right\rfloor,$$

and

$$(17) \quad \deg\{P(x)\} \geq \lambda_{n,k}.$$

PROOF: By induction on n . The result is true for $n \leq k$ because $\lambda_{n,k} = 0$ for $0 \leq n \leq k$. Suppose now that $n \geq k + 1$, and that the lemma holds for n replaced by any smaller number.

From the representation (12), and the induction hypothesis with n replaced by $n - k$,

$$\ell_{k+1} \geq 2 \left\lfloor \frac{n-k-1}{k} \right\rfloor.$$

From Lemma 2,

$$\ell_1 \geq 2 + \ell_{k+1} \geq 2 \left\lfloor \frac{n-1}{k} \right\rfloor$$

and (16) is proved. For (17), we use the representation (11) and the induction hypothesis with n replaced by $n - 1$. Then

$$\deg\{P(\mathbf{x})\} \geq \ell_1 + \lambda_{n-1,k} \geq \lambda_{n,k}. \quad \square$$

LEMMA 4. Suppose that $P(\mathbf{x})$ is a polynomial which has character $H_{n,k}(\ell_1, \ell_2, \dots, \ell_n)$ and with minimal total degree

$$d_{n,k} = \ell_1 + \ell_2 + \dots + \ell_n.$$

Then

$$(18) \quad d_{n,k} = \lambda_{n,k},$$

and

$$(19) \quad \ell_1 = 2 \left\lfloor \frac{n-1}{k} \right\rfloor.$$

PROOF: Result (18) follows from the fact that (10) shows that equality is possible in (17), using the generalised discriminant. From (11), (17) and (18), we have $\ell_1 = d_{n,k} - \deg\{Q\} \leq \lambda_{n,k} - \lambda_{n-1,k}$. But from (16),

$$\ell_1 \geq 2 \left\lfloor \frac{n-1}{k} \right\rfloor = \lambda_{n,k} - \lambda_{n-1,k}$$

and therefore

$$\ell_1 = \lambda_{n,k} - \lambda_{n-1,k} = 2 \left\lfloor \frac{n-1}{k} \right\rfloor. \quad \square$$

These preliminaries are the principal requirements for our main theorem:

THEOREM 1. The generalised discriminant theorem. *The non-trivial symmetric polynomial $P(x_1, x_2, \dots, x_n)$ of least total degree (with coefficients in a field of characteristic zero) which vanishes identically when any $k + 1$ of its variables x_i are equal is unique up to an overall scale factor, and it is called the generalised discriminant $P_{n,k}(\mathbf{x})$. It may be expressed as*

$$(20) \quad P_{n,k}(\mathbf{x}) = \sum_{\sigma \in S_n} \sigma \left\{ \prod_{\ell=1}^k \prod_{\substack{i < j \\ i, j \in I_\ell}} (x_i - x_j)^2 \right\}$$

where σ denotes any permutation of the indices $1, 2, \dots, n$. These indices have been grouped into disjoint sets I_1, I_2, \dots, I_k . The sets I_1 , then I_2 , then I_3, \dots ,

then I_r each consist of $q + 1$ indices taken sequentially from $1, 2, \dots, r(q + 1)$ and $I_{r+1}, I_{r+2}, \dots, I_k$ consist of q indices similarly taken from $r(q + 1) + 1, r(q + 1) + 2, \dots, n$. Here $q = \lfloor n/k \rfloor$ and $r = n \bmod k$.

The minimal degree of $P(\mathbf{x})$ is

$$(21) \quad \deg\{P_{n,k}(\mathbf{x})\} = kq(q - 1) + 2rq.$$

PROOF: The representation (20) is the same as (4), except that the indices are used in natural order for descriptive simplicity. The result (21) follows from (5) and (18), and the observation that (21) is trivial for $n \leq k$, because both $P(\mathbf{x})$ and $P_{n,k}(\mathbf{x})$ are nonzero constants (they contain too few variables to make equal).

The rest of the proof is by induction on n . From (11), (19) and the induction hypothesis,

$$P(\mathbf{x}) = \alpha x_1^{2\lfloor (n-1)/k \rfloor} P_{n-1,k}(x_2, x_3, \dots, x_n) + \text{lower order terms in } x_1,$$

for some non-zero constant α . From (4), (8) and (9),

$$P_{n,k}(\mathbf{x}) = \beta x_1^{2\lfloor (n-1)/k \rfloor} P_{n-1,k}(x_2, x_3, \dots, x_n) + \text{lower order terms in } x_1,$$

for some non-zero constant β . Therefore $S(\mathbf{x}) := \alpha P_{n,k}(\mathbf{x}) - \beta P(\mathbf{x})$ is a polynomial of total degree $\lambda_{n,k}$ and some character $H_{n,k}(\ell'_1, \ell'_2, \dots, \ell'_n)$ but degree less than $2\lfloor (n - 1)/k \rfloor$ in x_1 . This would contradict Lemma 4, unless $S(\mathbf{x}) \equiv 0$. \square

CONCLUSION

The generalised discriminant $P_{n,k}(\mathbf{x})$ does not appear to have been considered previously despite its fairly natural appearance. In the case $k = 1$, any symmetric polynomial $T(\mathbf{x})$ in the variables x_1, x_2, \dots, x_n which vanishes when any two of these variables are equal must be divisible by the ordinary discriminant $P_{n,1}(x)$ [4]. The corresponding result for the generalised discriminant $P_{n,k}(\mathbf{x})$ is not true when $k > 1$, and this fact presumably accounts for its lower profile. A similar phenomenon arises for certain ‘generalised resultants’ which were constructed by Pragacz [5]; these turn out to be super Schur functions [3].

Generalised discriminants were originally introduced to solve the problem of how the formula for the denominator polynomial of a vector-valued Padé approximant reduces in the scalar case [1]; the solution was given by Woodcock and Graves-Morris [7] but assuming the results of this paper.

REFERENCES

- [1] P.R. Graves-Morris, 'Problem 92-20', *SIAM Review* **34** (1992), 649-650.
- [2] S. Lang, *Algebra*, Chapter 5 Section 9 (Addison-Wesley, Reading, Mass., 1967).
- [3] I.G. Macdonald, *Symmetric functions and Hall polynomials* (Oxford University Press, Oxford, 1979).
- [4] M. Marcus, *Introduction to modern algebra* (Dekker, New York, 1978).
- [5] P. Pragacz, 'A note on the elimination theory', *Nederl. Akad. Wetensch. Indag. Math.* **49** (1987), 215-221.
- [6] R.J. Wilson, *Introduction to graph theory* (Longman, London, 1985).
- [7] C.F. Woodcock and P.R. Graves-Morris, 'Solution to Problem 92-20', *SIAM Review* **35** (1993), 649-651.

Institute of Mathematics and Statistics
University of Kent
Canterbury
Kent CT2 7NF
England

Department of Mathematics and Statistics
University of Canterbury
Private Bag 4800
Christchurch
New Zealand