

### CORRIGENDUM

## Variation in activity levels amongst dogs of different breeds: results of a large online survey of dog owners from the UK – CORRIGENDUM

Emily Pickup<sup>1</sup>, Alexander J. German<sup>1,2</sup>, Emily Blackwell<sup>3</sup>, Mark Evans<sup>4</sup> and Carri Westgarth<sup>1,5</sup>

<sup>1</sup>*Institute of Veterinary Science, University of Liverpool, Neston, UK*

<sup>2</sup>*Institute of Ageing and Chronic Disease, University of Liverpool, Neston, UK*

<sup>3</sup>*School of Clinical Veterinary Science, University of Bristol, Langford, UK*

<sup>4</sup>*Independent Veterinary Consultant, Guildford, UK*

<sup>5</sup>*Institute of Infection and Global Health, University of Liverpool, Liverpool, UK*

*Journal of Nutritional Science* (2021), vol. 10, e93, page 1 of 1

doi:[10.1017/jns.2017.7](https://doi.org/10.1017/jns.2017.7). Published by Cambridge University Press, 17 April 2017

This article was published with errors in the first sentence of the section ‘Off-lead exercise’ on page 5. The correct sentence is below.

#### *Off-lead exercise*

Some breeds were **less** commonly allowed off the lead in public ( $P < 0.001$ ; Table 1), including: chow chow (8, 73% **not** let off the lead); Siberian husky (67, 61%); Pyrenean mountain dog (3, 60%); Alaskan Malamute (31, 60%); and St Bernard (5, 56%).

The authors apologise for the error.

#### Reference

Pickup, E., German, A., Blackwell, E., Evans, M., & Westgarth, C. (2017). Variation in activity levels amongst dogs of different breeds: Results of a large online survey of dog owners from the UK. *Journal of Nutritional Science*, **6**, E10. doi:[10.1017/jns.2017.7](https://doi.org/10.1017/jns.2017.7)