
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editors, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
395A Hatfield Road,
St Albans, Herts AL4 0XQ.
England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

ASHFORD, R. W. & BARNISH, G. (1989) *Strongyloides fuelleborni* and similar parasites in animals and man. In: *Strongyloidiasis: a major roundworm infection of man* (editor, D. I. Grove) pp. 271–286. Taylor & Francis: London.

GARSIDE, P. & BEHNKE, J. M. (1989) Primary infection of hamsters with *Ancylostoma ceylanicum*. *Parasitology*, **98**, 283–289.

Contents

| | Pages |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| Studies of the relationships between <i>Schistosoma nasale</i> and <i>S. spindale</i> and their snail host <i>Indoplanorbis exustus</i> . J. DE BONT, J. VERCRUYSSSE, D. VAN AKEN, V. R. SOUTHAGTE and D. ROLLINSON | 1-7 |
| Cercarial shedding patterns of <i>Schistosoma bovis</i> and <i>S. haematobium</i> from single and mixed infections of <i>Bulinus truncatus</i> . A. MOUAHID, H. MONÉ, A. CHAIB and A. THÉRON | 8-14 |
| Natural infection of <i>Lymnaea truncatula</i> by the liver fluke <i>Fasciola hepatica</i> in the Porma Nasin, León, NW Spain. Y. MANGA-GONZALEZ, C. GONZALEZ-LANZA and C. B. OTERO-MERINO | 15-27 |
| <i>Diplostomum spathaceum</i> (Rud. 1819): effect of miracidial age and lifespan on miracidial infectivity. G. D. B. WAADU | 28-30 |
| The effects of age, sex and breed of chicken on susceptibility to <i>Clinostomum complanatum</i> (Digenea: Clinostomatidae). A. M. N. KALANTAN, M. ARFIN and W. A. NIZAMI | 31-37 |
| The route of infection of <i>Paragonimus westermani</i> (diploid type) cercariae in the freshwater crab, <i>Geothelphusa dehaani</i> . T. SHIBAHARA | 38-42 |
| Studies on the life-cycle of <i>Pleurogenoides ovatus</i> Rao, 1977 (Trematoda: Pleurogenetinae). K. P. JANARDANAN and P. K. PRASADAN | 43-50 |
| Characterization of the <i>Opisthorchis viverrini</i> genome. R. SERMSWAN, S. MONGKOLSUK and S. SIRISINHA (Research Note) | 51-54 |
| Release and survival of <i>Echinococcus</i> eggs in different environments in Turkana, and their possible impact on the incidence of hydatidosis in man and livestock. T. M. WACHIRA, C. N. L. MACPHERSON and J. M. GATHUMA | 55-61 |
| A comparative study of the protein content of some helminths and the suitability of assay methods. S. M. A. ABIDI and W. A. NIZAMI | 62-66 |
| Identification of human hookworm in failed-treatment cases using Chinese hamsters (<i>Cricetulus griseus</i>) and scanning electron microscopy. A. C. EVANS, T. J. M. DALY and M. B. MARKUS | 67-72 |
| Secretion and excretion of endogenously synthesized lipids by adult <i>Haemonchus contortus</i> <i>in vitro</i> . J. KAPUR and M. L. SOOD (Research Note) | 73-75 |
| Periodicity of <i>Dirofilaria immitis</i> microfilariae in a dog from Muheza district, Tanzania. Y. G. MATOLA (Research Note) | 76-78 |