

A NOTE ON A RESULT OF MAHLER'S

INDAR S. LUTHAR

(Received 4 November 1965)

In a recent paper, Mahler [2] proved that for any algebraic number field K of degree n and discriminant d there exists a constant C depending only on n and d such that for any ceiling $\lambda(\mathfrak{p})$ of K there exists a basis $\alpha_1, \dots, \alpha_n$ of the corresponding ideal \mathfrak{a}_λ such that

$$\begin{aligned} C^{-(n-1)}\lambda(\mathfrak{q}) &\leq |\alpha_k|_{\mathfrak{q}} \leq C\lambda(\mathfrak{q}) \quad \text{for all } \mathfrak{q}, \\ C^{-n}\lambda(\mathfrak{r}) &\leq |\alpha_k|_{\mathfrak{r}} \leq \lambda(\mathfrak{r}) \quad \text{for all } \mathfrak{r}; \\ &(k = 1, 2, \dots, n). \end{aligned}$$

Moreover, if \mathfrak{p}_r denotes the rational prime below r , then

$$|\alpha_k|_{\mathfrak{r}} = \lambda(\mathfrak{r})$$

for all r for which $\mathfrak{p}_r > C^n$.

For notations and definitions we refer the reader to the above-mentioned paper of Mahler.

The object of this note is to prove the same result with a constant C , which differs from Mahler's constant by a factor which approaches zero exponentially as n approaches infinity. We remark that for small n , Mahler's constant is better than ours.

Let β_1, \dots, β_n be a basis of \mathfrak{a}_λ and let

$$F(x) = F(x_1, \dots, x_n) = \sum_{\mathfrak{q}} \lambda(\mathfrak{q})^{-1} |\beta_1 x_1 + \dots + \beta_n x_n|_{\mathfrak{q}}.$$

Then $F(x)$ is a symmetric convex distance function and the volume V of the convex body

$$F(x) \leq 1$$

is given by the formula:

$$V = \frac{2^n \pi^{r_s}}{n! |\sqrt{d}|}.$$

We now use a theorem of Mahler [1] and Hermann Weyl [3] to obtain a unimodular matrix (g_{ij}) such that

$$\prod_{h=1}^n F(g_{h1}, \dots, g_{hn}) \leq \frac{\gamma_n}{V}$$

where γ_n is a certain constant depending only on n . If we write

$$(1) \quad \begin{cases} \alpha_1 = g_{11}\beta_1 + \dots + g_{1n}\beta_n \\ \alpha_n = g_{n1}\beta_1 + \dots + g_{nn}\beta_n \end{cases}$$

and

$$(2) \quad m_h = F(g_{h1}, \dots, g_{hn}) = \sum_q \lambda(q)^{-1} |\alpha_h|_q$$

then $\alpha_1, \dots, \alpha_n$ is a basis of \mathfrak{a}_λ and

$$(3) \quad \prod_{h=1}^n m_h \leq \gamma_n V^{-1} = n! \gamma_n 2^{-n} \pi^{-r_s} |\sqrt{d}|.$$

By the inequality on arithmetic and geometric means we get

$$\frac{1}{n} m_h = \frac{1}{n} \sum_q \lambda(q)^{-1} |\alpha_h|_q \geq [4^{-r_s} \prod_q \lambda(q)^{-n(q)} \cdot |N(\alpha_h)|]^{1/n}$$

i.e.,

$$N(\mathfrak{a}_\lambda)^{-1} |N(\alpha_h)| \leq 4^{r_s} \left(\frac{m_h}{n}\right)^n.$$

Since $\alpha_h \in \mathfrak{a}_\lambda$, therefore $N(\mathfrak{a}_\lambda) \leq |N(\alpha_h)|$ and the above inequality gives

$$(4) \quad m_h \geq n \cdot 4^{-r_s/n}.$$

Using (2), (3), and (4) we obtain

$$|\alpha_h|_q \leq \lambda(q) m_h = \lambda(q) \frac{\prod_{\substack{i=1 \\ i \neq h}}^n m_i}{\prod_{\substack{i=1 \\ i \neq h}}^n m_i} \leq \lambda(q) C$$

with

$$C = n! \gamma_n n^{1-n} \pi^{-r_s} 2^{-(r_1 + (2r_s/n))} |\sqrt{d}|.$$

The remaining assertions of Mahler's theorem with this C now follow from his lemmas 1 and 2.

We remark finally that the ratio of Mahler's constant to the constant obtained here is

$$\frac{2^{-r_s/n} n^{\frac{1}{2}(1-n)} \gamma_n \pi^{-\frac{1}{2}n} \Gamma\left(\frac{n}{2} + 1\right) |\sqrt{d}|}{n! \gamma_n n^{1-n} \pi^{-r_s} 2^{-(r_1 + 2r_s/n)} |\sqrt{d}|} \sim \frac{\sqrt{n}}{2\sqrt{2}} 2^{r_s/n} \left(\frac{4}{\pi}\right)^{\frac{1}{2}r_1} \left(\frac{e}{2}\right)^{\frac{1}{2}n}.$$

References

- [1] K. Mahler, *Acta Mathematica* **68** (1937), 109–144.
- [2] K. Mahler, *J. Austr. Math. Soc.* **4** (1964), 425–448.
- [3] H. Weyl, *Proc. London Math. Soc.* **47** (1942), 268–289.

University of Illinois
Urbana, Illinois .