

---

## INSTRUCTION TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editor, Journal of Helminthology,  
London School of Hygiene and Tropical Medicine,  
Winches Farm Field Station,  
395 Hatfield Road,  
St Albans, Herts AL4 0XQ,  
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0 × 20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0 × 20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

67

# Contents

|  | Pages              |
|--|--------------------|
| Experimental <i>Schistosoma mansoni</i> infection in vervet monkeys ( <i>Cercopithecus aethiops</i> ) in Kenya: I. Susceptibility to a primary infection. R. F. STURROCK, F. M. OTIENO, R. TARARA, R. KIMANI, R. HARRISON and J. G. ELSE | 79-92              |
| Scanning electron microscope studies of the mucosa of rats infected with <i>Hymenolepis diminuta</i> (Cestoda). JEAN MARTIN and CELIA HOLLAND  | 93-99              |
| A comparative study on esterases from three species of <i>Raillietina</i> . M. P. BALASUBRAMANIAN, S. DHANDAYUTHAPANI, K. NELLAIAPPAN and K. RAMALINGAM  | 101-105            |
| The relevance of <i>in vitro</i> anthelmintic screening tests employing the free-living stages of trichostrongylid nematodes. O. F. IBARRA and D. C. JENKINS   | 107-112            |
| Trematodiasis in Turkey: comparative efficacy of triclabendazole and niclofolan against natural infections of <i>Fasciola hepatica</i> and <i>F. gigantica</i> in sheep. N. GÜRALP and R. TINAR  | 113-116            |
| Microfilarial periodicity in <i>Mastomys natalensis</i> . A. B. REDDY, R. CHANDRASHEKAR, U. R. RAO and D. SUBRAHMANYAM   | 117-121            |
| The relationship between the size of <i>Lymnaea truncatula</i> naturally infected with <i>Fasciola hepatica</i> and the intensity and maturity of the redial infection. G. SMITH   | 123-127            |
| <i>Litomosoides carinii</i> in jirds ( <i>Meriones unguiculatus</i> ): ability to retard development of challenge larvae can be transferred with cells and serum. D. J. WEINER, D. ABRAHAM and R. D'ANTONIO                              | 129-137            |
| <i>Toxocara canis</i> infection in dogs in Beersheba, Israel. (Research Note). E. M. GROSS, R. ZEITAN and V. TOROK   | 139-141            |
| Induction of peripheral blood eosinophilia in mice by excretory and secretory antigen of <i>Toxocara canis</i> larvae. K. SUGANE and T. OSHIMA   | 143-147            |
| An SEM study of the cephalic region, buccal cavity and male tail of the species of the genus <i>Strongylus</i> Müller, 1780 (Nematoda, Strongyloidea). LYNDIA M. GIBBONS   | 149-158            |
| Studies on infectivity, longevity and fecundity of <i>Ancylostoma ceylanicum</i> in golden hamsters. P. K. S. VISEN, J. C. KATIYAR and A. B. SEN   | 159-163            |
| The histopathology of <i>Neoechinorhynchus hutchinsoni</i> Datta, 1936 (Neoacanthocephala: Neoechinorhynchidae) infection in <i>Nemacheilus kashmirensis</i> Hora. M. K. RAINA and P. L. KOUL  | 165-168            |
| Histochemical localization of phosphomonoesterases in <i>Avitellina lahorea</i> Woodland, 1927 (Cestoda: Anoplocephalida). R. FAROOQ and H. U. FAROOQI   | 169-173            |
| Announcements  | 122, 138, 142, 174 |
| Corrigenda   | 100                |