

Conclusions. The INTEGRATE-HTA methodology helped to identify interactions between domains and significantly expanded the perspective on a technology. It improved patient-centered understanding and facilitated the discussion of single aspects.

PP78 Analysis Of Resource Utilization In Psoriasis Care In The Brazilian Health System

Larissa de Araujo Costa Andrade (larissa.andrade@abbvie.com), Ricardo Moreira and Vinicius Vitale

Introduction. Psoriasis is associated with metabolic disorders and has a significant negative impact on patients' quality of life. There is little information about treatment patterns and the resources used to treat psoriasis and its associated comorbidities. This study aimed to evaluate the frequency of metabolic syndrome (MetS) and estimate the costs associated with psoriasis care in the Brazilian health system.

Methods. A cross-sectional observational study was conducted of 293 Brazilian patients with psoriasis who attended selected dermatology outpatient centers in Brazil during a one-year period. Patients underwent detailed skin and rheumatologic assessments and a series of laboratory tests. The annual costs were estimated by multiplying the amount of each resource consumed by its unit cost.

Results. The prevalence rate of MetS was high in this cohort (50%). Other metabolic disorders were more common in this group than in the general population, including hypertension (62%), diabetes (31%), dyslipidemia (75%), and obesity (53%). The disease duration was, on average, 17.2 years. There were 2,713 consultations reported by 288 patients (average of nine per patient per year). The most common specialists consulted were: dermatologists (93%); general practitioners (41%); cardiologists (27%); rheumatologists (18%); and endocrinologists (13%). There were 110 non-physician visits (average of 11 per patient per year), of which the most frequently consulted specialists were nutritionists (26%), psychologists (24%), and nurses (22%). A total of 279 patients (95%) had at least one medical test (average of 18 per patient per year). Among the treatments required for other conditions, 65% (n = 149) of prescribed medicines were for the cardiovascular system and 24% (n = 5/21) of hospitalizations were due to diseases of the circulatory system; myocardial revascularization was the costliest procedure (USD 2,298).

Conclusions. MetS is frequently associated with psoriasis and directly affects health resource utilization in the Brazilian health system by increasing consultation costs. Further research is needed to evaluate the impact of a multidisciplinary treatment approach for psoriasis patients with MetS.

PP79 Impact Of Hidradenitis Suppurativa On Healthcare Resource Utilization

Larissa de Araujo Costa Andrade (larissa.andrade@abbvie.com), Ricardo Moreira and Vinicius Vitale

Introduction. Hidradenitis suppurativa (HS) is a debilitating, chronic inflammatory skin disease characterized by painful nodules and abscesses. HS has a strong impact on patient quality of life. In Brazil, the prevalence of HS is estimated at 0.4 percent. Medical and surgical treatments have low effectiveness and disease recurrence is common, which affects health system costs. This study aimed to assess how HS patients utilize medical care (emergency and inpatient care) in Brazil and to describe the all-cause costs.

Methods. Data were retrieved from a public healthcare claims database (DATASUS), which provides access to information regarding health services and costs. Data from DATASUS were used to perform a cost-identification analysis on patients with HS who used health services over a two-year period. A retrospective bottom-up approach was used to estimate direct costs, multiplying the amount of each medical resource consumed by its unit cost.

Results. Over the two-year period, 90 patients (16%) with HS received inpatient care (151 procedures) at a total cost of BRL 83,520 (USD 21,715). Surgeries were the most frequently performed (73% of total) and expensive procedures, costing BRL 73,122 (USD 19,011; 88% of total costs), followed by clinical treatments (BRL 8,354 [USD 2,172]; 10%), and physician consulting (BRL 1,659 [USD 431]; 2%). For the 500 patients treated in the emergency department (total cost BRL 3,027 [USD 787]), the most frequently received services were physician consulting (34%), nursing care (12%), and minor surgeries (11%). Each patient received, on average, three procedures over the two-year period.

Conclusions. HS is a high-burden disease, as demonstrated by the high healthcare resource utilization among patients. Since DATASUS is a public database, the costs presented reflect a government reference price and do not consider local costs, which is a limitation of this study. Health managers should be aware of this finding, although further research is needed to investigate the effect of healthcare utilization on patient outcomes.

PP80 A Systematic Review Of The Gugging Swallowing Screen For Assessing Dysphagia

Seon Heui Lee (sunarea87@gachon.ac.kr)

Introduction. Dysphagia is a clinical burden that can lead to serious complications like aspiration and pneumonia. Complications often result in longer hospital stays or an increased mortality rates. The Gugging Swallowing Screen (GUSS) assesses swallowing ability in patients by allowing separate evaluations for non-fluid and fluid textures, and is a potentially useful tool for determining the risk of aspiration and dysphagia. The purpose of this study is to analyze the validity and effectiveness of GUSS for dysphagia screening.

Methods. We conducted a systematic review by searching the following electronic databases: Medline, EMBASE, The Cochrane Library, KoreaMed, the Research Information Sharing Service, and the Korean Studies Information Service System. We included studies related to dysphagia screening with GUSS that were published in English or Korean up to November 2018.

Results. Of 297 unique studies identified, 219 were reviewed by two independent reviewers. Finally, eight articles were identified as being relevant for this study. With regard to validity, GUSS had a sensitivity ranging from 90 to 100 percent and a specificity of between 50 and 88 percent. In addition, GUSS results significantly correlated with the results of the videofluoroscopic swallow study and the fiberoptic endoscopic evaluation of swallowing. In terms of effectiveness, early systematic dysphagia screening with GUSS by nurses reduced the duration of screening and rate of pneumonia, compared with the control group ($p = 0.004$). The incidence of X-ray verified pneumonia in the GUSS group was also significantly lower than in the clinical screening group ($p < 0.01$), although there was no difference in the occurrence of pneumonia, compared with the 10 mL water swallowing test.

Conclusions. Results showed that GUSS is a reliable and sensitive tool for screening patients for dysphagia. This early and systematic assessment can reduce the occurrence of aspiration and pneumonia, although further research is needed to establish the effectiveness of GUSS.

PP81 Real World Data: The Early Access To Medicines Scheme Catches The Worm

Adam Hall, Lok Wan Liu (lokwan.liu@PAREXEL.com), Richard Macaulay and Sean Walsh

Introduction. The Early Access to Medicines Scheme (EAMS) aims to provide access to medicines prior to market authorization for patients with severe, life-threatening diseases who do not have adequate treatment options. An EAMS designation enables the potential collection of United Kingdom-specific real world evidence (RWE) prior to health technology assessment (HTA) by the National Institute for Health and Care Excellence (NICE). This research evaluates whether RWE is being gathered through the EAMS and utilized to support HTA submissions.

Methods. All EAMS designations as of 7 November 2018 were identified from the Medicines and Healthcare products Regulatory Agency website. For products with final NICE guidance, all publicly-available NICE documentation was reviewed.

Results. Sixteen product and indication pairings with an EAMS designation were identified, with 12 having received final NICE guidance (11 were recommended, 3 were recommended for temporary reimbursement via the Cancer Drugs Fund, and 2 were not recommended). Of the 11 recommended products, seven had references to the number of patients or sites with product access through the EAMS, but only one (dupilumab for atopic dermatitis) had detailed data collected during the EAMS period. The manufacturer of dupilumab reported baseline demographics and disease characteristics from a cohort of 35 patients treated under the EAMS to inform the generalizability of trial populations for clinical practice. Follow-up results from this cohort demonstrated that real-world data on dupilumab effectiveness was comparable with the clinical trial data, despite a higher proportion of patients in the real-world cohort receiving immunosuppressant therapy, which makes improvements in efficacy harder to achieve. The committee also noted that the RWE presented supported the understanding of dupilumab's long-term clinical effectiveness and informed assumptions for the economic model.

Conclusions. To date, the majority of products receiving an EAMS designation have not presented RWE at NICE reappraisal. The case of dupilumab illustrated how RWE collected through the EAMS can be used to reduce uncertainty around how clinical trial data can be translated into clinical practice. In the future, RWE may increasingly be used to help inform NICE decisions.

PP83 A Conceptual Decision-Making Framework For Pharmaceutical Innovations

Cornelis Boersma (c.boersma01@umcg.nl), Joost Geenen and Maarten Postma

Introduction. The trend of growing healthcare expenditures is unsustainable in many countries. The increasing pressure on healthcare budgets due to, for example, population ageing, increasing numbers of patients with chronic diseases (including multimorbidity), and the introduction of new pharmaceutical innovations, leads to political and societal debate. In particular, the introduction of expensive pharmaceutical innovations causes a lot of discussion and uncovers various paradoxes and dilemmas. There is a societal demand for innovation focused on existing medical needs (e.g., oncological, immune-mediated inflammatory, and orphan diseases), but the price of pharmaceutical innovations is a barrier to patient access. As a consequence, systems try to introduce measures or incentivize market forces to improve access for patients, while also containing budget impact. This does not always lead to better access and affordability. The aim of this study was to develop and test a conceptual decision-making framework for pharmaceutical innovations.

Methods. A retrospective study was conducted to identify the successes and challenges of decision-making systems across Europe. A conceptual decision-making framework, including proposed procedures, criteria, and health technology assessment (HTA) requirements (including tools), was developed and tested based on specific case examples (e.g. oncology and hepatitis C).

Results. The conceptual decision-making framework comprised an algorithm for relevant decision-making criteria (e.g. clinical evidence, medical need, cost-effectiveness, and budget impact). The algorithm was developed hierarchically and ranked the criteria in order to optimally inform various types of investment decisions. This novel approach to conducting budget impact analyses resulted in more realistic predictions of the burden of pharmaceutical innovations on healthcare budgets, and can be used as part of horizon-scanning processes to inform healthcare decision making. Results from selected case examples are presented.

Conclusions. The conceptual decision-making framework and proposed method for budget impact predictions will allow for more balanced future healthcare investment decisions.

PP84 Different Interpretation Of Evidence By A Health Technology Assessment Body And A Decision Maker

Charalabos-Markos Dintsios (dintsios@hhu.de), Jörg Ruof, Franziska Worm and Matthias Herpers