



Letter

Cite this article: Simkins LM, Greenwood SL, Winsborrow MCM, Bjarnadóttir LR, Lepp AP (2022). Advances in understanding subglacial meltwater drainage from past ice sheets. *Annals of Glaciology* 63(87-89), 83–87. <https://doi.org/10.1017/aog.2023.16>

Received: 19 October 2022
Revised: 24 January 2023
Accepted: 25 February 2023
First published online: 17 April 2023


Keywords:

Geomorphology; glacier hydrology; sedimentology

Author for correspondence:

Lauren M. Simkins,
E-mail: lsimkins@virginia.edu

Advances in understanding subglacial meltwater drainage from past ice sheets

Lauren M. Simkins¹ , Sarah L. Greenwood², Monica C. M. Winsborrow³, Lilja R. Bjarnadóttir⁴ and Allison P. Lepp¹

¹Department of Environmental Sciences, University of Virginia, Charlottesville, Virginia, USA; ²Department of Geological Sciences, Stockholm University, 10691 Stockholm, Sweden; ³Department of Geosciences, Centre for Arctic Gas Hydrate, Environment and Climate, UiT The Arctic University of Norway, 9037 Tromsø, Norway and ⁴Geological Survey of Norway, 7491 Trondheim, Norway

Abstract

Meltwater drainage beneath ice sheets is a fundamental consideration for understanding ice–bed conditions and bed-modulated ice flow, with potential impacts on terminus behavior and ice-shelf mass balance. While contemporary observations reveal the presence of basal water movement in the subglacial environment and inferred styles of drainage, the geological record of former ice sheets, including sediments and landforms on land and the seafloor, aids in understanding the spatiotemporal evolution of efficient and inefficient drainage systems and their impact on ice-sheet behavior. We highlight the past decade of advances in geological studies that focus on providing process-based information on subglacial hydrology of ice sheets, how these studies inform theory, numerical models and contemporary observations, and address the needs for future research.

1. Introduction

Liquid water beneath ice sheets influences ice-flow organization and velocity (Kyrke-Smith and others, 2014; Bell and others, 2007; Larter and others, 2019), subglacial sediment rheology and transport (Damsgaard and others, 2020; Minchew and Meyer, 2020), grounding-line behavior (Horgan and others, 2013; Fried and others, 2015) and ice-shelf mass balance (Le Brocq and others, 2013; Alley and others, 2016). Ice-sheet response, however, is contingent on subglacial water supply and drainage organization (Röthlisberger, 1972; Walder, 1986; Schoof, 2010). While some components of subglacial hydrological systems are relatively stable (i.e. fixed), such as large subglacial lakes beneath the East Antarctic ice sheet (Kapitsa and others, 1996) and incised bedrock channels (Kirkham and others, 2020), other reservoirs and drainage pathways are more transient and evolve through time and space with non-linear and spatially heterogeneous impacts on ice-sheet behavior (Schroeder and others, 2013; Andrews and others, 2014; Hoffman and others, 2016; Siegfried and others, 2016; Rada and Schoof, 2018). Beyond the grounding line, contemporary sediment plumes emanating from marine-terminating outlet glaciers of the Greenland ice sheet observed via satellite imagery (Fried and others, 2015; Schild and others, 2016) and surficial expressions of channelization beneath Antarctic ice shelves (Le Brocq and others, 2013; Alley and others, 2016) indicate active subglacial hydrological systems upstream.

Major advances in observing contemporary ice-sheet hydrology, such as radar specularities (Schroeder and others, 2013) and repeat satellite measurements (Fricker and others, 2016), reveal spatiotemporal evolution of basal water transmission on sub-decadal scales. Yet, the limited nature of long-term (decadal to millennial) observations impedes holistic perspectives on the modes and magnitudes of water drainage beneath ice sheets and their consequences for ice-sheet behavior. In formerly glaciated landscapes and continental margins, relict subglacial water drainage is recorded by meltwater landforms and sedimentological successions (Fig. 1; Kehew and others, 2012; Lee and others, 2015; Greenwood and others, 2016; Esteves and others, 2017). Channels (broadly defined) incised into bedrock and sediments and positive-relief esker ridges record drainage styles and organization (Storarr and others, 2014; Zoet and others, 2019; Lewington and others, 2020) and, in some cases, associated ice-margin retreat behavior (Livingstone and others, 2020; Simkins and others, 2021) and creation of tidal embayments in grounding lines (Horgan and others, 2013). Distinct meltwater plume deposits and hydrologically sorted sediments reveal relative magnitudes and frequency of water drainage into the ocean, precursory, synchronous or resulting glacial environment changes, and geochemical signatures of sediment and water provenance (Witus and others, 2014; O'Regan and others, 2021; Lepp and others, 2022). Such empirical observations based on the geological record have long guided and continue to inform, and challenge, glaciological theory (e.g. Walder and Hallet, 1979; Boulton and others, 2009; Hewitt, 2011).

2. Recent scientific advances from paleo-ice sheets

Over the past decade, advances in geophysical methods, growing data accessibility and sedimentological studies from deglaciated terrains have pushed the boundaries of subglacial






Fig. 1. (a) Murtoo pathway within glacially streamlined terrain in central Finland (Mäkinen and others, 2017; Ojala and others, 2019). Data: LiDAR-based DEM from the National Land Survey of Finland. (b) Meltwater channels and eskers drape and incise drumlins in the Bothnian Sea (Greenwood and others, 2017). Data: MBES-based DEM from the Swedish Maritime Administration. (c) Meltwater channel incised retreat moraines (red dots) on Thor Iversenbanken in the Central Barents Sea (Esteves and others, 2017). Data: MAREANO MBES-based bathymetry from the Norwegian Mapping Authority. (d) Meltwater corridor in which channels cross-cut grounding zone wedges (red dots) in the Ross Sea, Antarctica (Simkins and others, 2021). Data: MBES-based DEM from cruise NBP15-02, available through the United States Antarctic Program Data Center. In (a)–(d), red dashed lines outline the encompassing areas of meltwater landforms. (e) CT scan and photograph of the upper 250 cm of sediment core NBP19-02 KC-08, collected in the Amundsen Sea, records meltwater plume events that emanated from the Thwaites Glacier grounding line (Lepp and others, 2022).

hydrology understanding and challenged concepts of the spatio-temporal evolution of water drainage beneath ice sheets. Near complete coverage of light detection and ranging (LiDAR) and satellite photogrammetry elevation data across terrestrial landscapes formerly glaciated by the European and North American ice sheets, give unprecedented views of paleo-subglacial meltwater landforms that range in relief from 10^{-1} to 10^2 m and lengths of 10^1 – 10^5 m, allowing holistic ice-sheet scale assessment of controls on drainage. While nowhere near as complete, increasing coverage and quality of bathymetry data from deglaciated continental shelves provide perspectives on water flow beneath marine-based ice sheets and implications for ice-sheet behavior at fine scales previously unseen. These offshore advances via multibeam echo sounding (MBES) surveys are facilitated by national hydrographic programs such as MAREANO (e.g. Esteves and others, 2017), marine geological repositories such as the Marine Geoscience Data System and researcher-led surveying and compilations (e.g. Greenwood and others, 2021). Additionally, 3-D seismic survey grids, albeit sparse but increasing in spatial coverage due to industry-academic relations, are unique datasets to assess temporal evolution of drainage pathways and internal architecture of meltwater landforms (e.g. Kirkham and others, 2021, 2022). These advances in terrain data acquisition and availability have not only permitted ice-sheet scale documentation of small-scale (meter to sub-meter) and intricate meltwater landforms, demonstrating their variability as well as near ubiquity, but have also uncovered both new types of landforms and little recognized meltwater landform assemblages, stimulating new hypotheses for meltwater landform genesis and understanding of the coupling to ice-flow and ice-margin behavior (Storror and others, 2014; Ojala and others, 2019; Kirkham and others, 2020).

Challenging the traditional binary categorization of subglacial drainage through either ‘channelized’ or ‘distributed’ pathways (Röthlisberger, 1972; Kamb, 1987), meltwater corridors found in the Northern Hemisphere (e.g. Peterson and others, 2018; Lewington and others, 2020), and on the Antarctic seafloor (Simkins and others, 2021) represent broad subglacial drainage pathways of meltwater landform assemblages that span 10^1 – 10^2 km in length. Corridor-like drainage systems have also been identified beneath the contemporary Greenland ice sheet (Hoffman and others, 2016; Davison and others, 2019) with complex configurations evolving depending on hydraulic gradients, meltwater input to the subglacial environment and sediment deformation (Davison and others, 2019). These corridors indicate co-existence of drainage styles, varying genetic erosional and depositional processes and waxing and waning of drainage magnitudes in time and space. Additionally, newly observed landforms in Scandinavia, termed murtoos, potentially bridge the long-standing gap in recognizing geomorphic evidence for distributed subglacial drainage (Mäkinen and others, 2017; Ojala and others, 2019). These low-relief triangular subglacial landforms oriented with their apex in the ice-flow direction (Fig. 1a) often occur with other meltwater landforms such as channels and eskers, and sediments which have undergone hydraulic sorting, ductile deformation and liquefaction (Becher and Johnson, 2021). Collectively, murtoo presence suggests efficient transitional drainage between channelized and distributed under high-pressure conditions, potentially in response to transient linked cavity-type drainage systems (Ojala and others, 2022) similar to those beneath the Greenland ice sheet (Hoffman and others, 2016). Both corridors and murtoos point to variable modes of drainage that co-exist or evolve in time and space including ‘efficient’ and ‘inefficient’ components, thus questioning the validity of assuming or parameterizing singular modes of subglacial water drainage.

A long-standing challenge in glacial geomorphology has been how to interpret the temporal significance of meltwater

landforms: the time required, and the stability of discharge required, for both landform and whole drainage pathway formation. Meltwater landform relations to other subglacial and ice-marginal landforms provide insights in this regard (e.g. Greenwood and others, 2017; Simkins and others, 2017; Ojala and others, 2019; Livingstone and others, 2020). For example, drumlins and mega-scale glacial lineations incised by channels and draped by eskers in the Bothnian Sea indicate a geomorphic switch from active bedform shaping to channelized water drainage overprinting stable bedforms, shortly before deglaciation (Fig. 1b; Greenwood and others, 2017). Here, interlinking channels and eskers of comparable sizes within a coherent drainage path highlight the transitory dominance of erosion and deposition in the subglacial environment. Episodic esker segment (‘bead’) deposition has long been inferred from the terrestrial landform-sediment record (De Geer, 1897; Banerjee and McDonald, 1975; Mäkinen, 2003). Livingstone and others (2020) demonstrate a tight relationship between esker beads and De Geer moraines in central Nunavut and infer time-transgressive landform building by drainage pathways to the ice margin. Similarly, meltwater channel incision through retreat moraines in the Barents Sea (Fig. 1c; Esteves and others, 2017) and variable incision of or draping by retreat moraines in the western Ross Sea (Simkins and others, 2017) indicate the relative persistence of channelized drainage during active ice-margin retreat. Embayments can form where meltwater channels drain at grounding lines (Horgan and others, 2013; Simkins and others, 2017) likely through grounding-line sediment non-deposition, with potential to enhance tidal action at and upstream of the grounding line as ocean water flushes in and out of the channel path as observed at the contemporary Whillans Ice Stream (Horgan and others, 2013). A corridor of over 80 meltwater channels on the Antarctic continental shelf (Fig. 1d; Simkins and others, 2021) had prolonged impacts on grounding-line behavior as larger magnitude grounding-line retreat events and grounding zone wedge deposition occurred while the channels within the corridor were active, compared to smaller retreat events and moraine deposition when the channels were inactive. While the mechanism for this relationship remains unknown, it possibly results from hydrological controls on sediment rheology and mobility that influence building of ice-marginal landforms that may or may not reduce effective water depths enough to counterbalance grounding-line buoyancy-driven retreat. Such observations of meltwater–grounding-line landform associations and the potential to document these over large tracts of paleo-ice-sheet beds offer new possibilities for constraining the time component of the meltwater landform record, as well as quantifying sediment loads and, for example, seasonal deposition of individual esker beads.

Complementary to geomorphological studies, sediment records from deglaciated continental shelves and proglacial lake basins elucidate the temporal persistence of subglacial and grounding-line water discharge and associated changes in ice-sheet configuration and behavior (e.g. Rüther and others, 2012; Lee and others, 2015; Avery and others, 2021; O’Regan and others, 2021; Lepp and others, 2022). Meltwater plume deposits offshore of Thwaites Glacier, Antarctica and Ryder Glacier, Greenland are a common feature associated with (or precursor to) glacier retreat and ice-shelf break up events, indicated by the millimeter-scale stratigraphy resolved by computed tomography (CT) scans and by grain-scale sedimentology (Fig. 1e; O’Regan and others, 2021; Lepp and others, 2022). Downcore stratigraphy and trace elemental ratios in cores that sample meltwater plume deposits reveal differences in relative magnitudes and frequencies of subglacial drainage into the ocean offshore of western and eastern Thwaites Glacier, and suggest greater magnitudes of sediment-laden water were delivered to the ocean in recent

centuries compared to the past several thousand years (Lepp and others, 2022). In the Baltic Sea basin, where proglacial varved sediments have long been used to document the pattern and pace of Fennoscandian ice-margin retreat, Avery and others (2021) find multi-decadal cycles of enhanced meltwater discharge through a 725 year varve series, ~15 000 years ago. In these paleo cases, and particularly where the former ice-sheet bed is now exposed, there is great potential for examining links between the temporal information archived in the distal sedimentological record of meltwater events and longevity of discharge, and the high-resolution geomorphology of the hydrological system responsible.

A recent body of work examining physical processes of and conditions for subglacial fluvial erosion, deposition and sediment mobility is an important step forward (Beaud and others, 2016, 2018; Damsgaard and others, 2017; Hewitt and Creyts, 2019; Kirkham and others, 2022; Stevens and others, 2022; V´erit´e and others, 2022). These studies build towards an integrated or continuum view of meltwater organization, depending on water supply and sources, basal conditions and substrate properties. Importantly, they make advances towards knowledge of where, over what timescales, and with what meltwater discharge regimes, sediments are mobilized and landforms may form, opening up the vast landform record to much more effective and accurate use as a document of coupled meltwater – ice flow – ice margin behavior in paleo-ice sheets over seasonal-to-millennial timescales.

3. Looking forward

To push the field of subglacial hydrology forward using the landform and sediment records of deglaciated regions, we need: increased geophysical data coverage in regions proximal to contemporary ice-sheet margins; coupled remote-sensing and field-based observations in terrestrial landscapes; reporting of quantitative sedimentologic and morphometric data and community building to work across disciplinary bounds and study-area silos. Of promise in narrowing knowledge gaps are emerging themes of research on deeper groundwater interactions with the ice–bed interface (Gustafson and others, 2022) and its implications for landform genesis (Boulton and others, 2009; Hermanowski and Piotrowski, 2019), understanding subglacial lake and ice-sheet surface connections to subglacial drainage systems (Greenwood and others, 2016; Simkins and others, 2017), assessment of the role of local (10^0 – 10^1 m relief) variability in bed conditions on drainage organization (Simkins and others, 2021) and the continued pursuit of constraining time for meltwater landform construction and evolution; each of these pursuits will benefit from more seamless integration of theory, numerical models and coupled geomorphological and chronological studies (Kirkham and others, 2022; Stevens and others, 2022). Additionally, higher-resolution topographic data from the surface of Mars offer opportunities to dig deeper into the surficial expressions of meltwater landforms (e.g. Butcher and others, 2020), whereby comparison with Earth’s meltwater landforms may offer new insights into key processes controlling their genesis. Here on Earth and on Mars, we need to be mindful of what we are not seeing, such as evidence for distributed, transient and hard-bed systems that might not leave their mark, in geomorphological and sedimentological records of subglacial hydrological systems. Additionally, when planning new research projects, those of us working on records of paleo-ice sheets and those studying contemporary ice sheets should draw on literature from the two respective fields to identify key gaps in understanding of subglacial hydrological forms and processes that will aid in assessing the future of the Greenland and Antarctic ice sheets.

Acknowledgments. The authors acknowledge their funding sources past, present and future that allow them to work toward better understanding subglacial processes; the traditional stewards of the global lands in which they work, some of whom directly interacted with extinct ice sheets and newly deglaciated landscapes following the Last Glacial Maximum; and those who collect and curate the data that support this work.

References

- Alley KE, Scambos TA, Siegfried MR and Fricker HA (2016) Impacts of warm water on Antarctic ice shelf stability through basal channel formation. *Nature Geoscience* **9**(4), 290–293.
- Andrews LC and 6 others (2014) Direct observations of evolving subglacial drainage beneath the Greenland ice sheet. *Nature* **514**(7520), 80–83.
- Avery RS and 6 others (2021) A 725-year integrated offshore terrestrial varve chronology for southeastern Sweden suggests rapid ice retreat ~15 ka BP. *Boreas* **50**(2), 477–496.
- Banerjee I and McDonald BC (1975) Nature of esker sedimentation. In Jopling AV and McDonald BC (eds), *Glaciofluvial and Glaciolacustrine Sedimentation*, vol. 23. SEPM Sp. Pub., pp. 132–154.
- Beaud F, Flowers GE and Venditti JG (2016) Efficacy of bedrock erosion by subglacial water flow. *Earth Surface Dynamics* **4**(1), 125–145.
- Beaud F, Flowers GE and Venditti JG (2018) Modeling sediment transport in ice-walled subglacial channels and its implications for esker formation and proglacial sediment yields. *Journal of Geophysical Research: Earth Surface* **123**(12), 3206–3227.
- Becher GP and Johnson MD (2021) Sedimentology and internal structure of murtoos-V-shaped landforms indicative of a dynamic subglacial hydrological system. *Geomorphology* **380**, 107644.
- Bell RE, Studinger M, Shuman CA, Fahnestock MA and Joughin I (2007) Large subglacial lakes in East Antarctica at the onset of fastflowing ice streams. *Nature* **445**(7130), 904–907.
- Bell RE, Banwell AF, Trusel LD and Kingslake J (2018) Antarctic surface hydrology and impacts on ice-sheet mass balance. *Nature Climate Change* **8**(12), 1044–1052.
- Boulton GS, Hagdorn M, Maillot PB and Zatsepin S (2009) Drainage beneath ice sheets: groundwater–channel coupling, and the origin of esker systems from former ice sheets. *Quaternary Science Reviews* **28**, 621–638.
- Butcher FE and 6 others (2020) Morphometry of a glacier-linked esker in NW Tempe Terra, Mars, and implications for sediment-discharge dynamics of subglacial drainage. *Earth and Planetary Science Letters* **542**, 116325.
- Damsgaard A and 5 others (2017) Sediment behavior controls equilibrium width of subglacial channels. *Journal of Glaciology* **63**(242), 1034–1048.
- Damsgaard A, Goren L and Suckale J (2020) Water pressure fluctuations control variability in sediment flux and slip dynamics beneath glaciers and ice streams. *Communications Earth & Environment* **1**(1), 1–8.
- Davison BJ, Sole AJ, Livingstone SJ, Cowton TR and Nienow PW (2019) The influence of hydrology on the dynamics of land-terminating sectors of the Greenland ice sheet. *Frontiers in Earth Science* **10**.
- De Geer G (1897) Om rullstensåsarnas bildningsätt. *GFF* **19**(5), 366–388.
- Esteves M, Bjarnadóttir LR, Winsborrow MC, Shackleton CS and Andreassen K (2017) Retreat patterns and dynamics of the Sentralbankrenna glacial system, Central Barents Sea. *Quaternary Science Reviews* **169**, 131–147.
- Fricker HA, Siegfried MR, Carter SP and Scambos TA (2016) A decade of progress in observing and modelling Antarctic subglacial water systems. *Philosophical Transactions of the Royal Society A* **374**(2059), 20140294.
- Fried MJ and 6 others (2015) Distributed subglacial discharge drives significant submarine melt at a Greenland tidewater glacier. *Geophysical Research Letters* **42**(21), 9328–9336.
- Greenwood SL, Clason CC, Helanow C and Margold M (2016) Theoretical, contemporary observational and palaeo-perspectives on ice sheet hydrology: processes and products. *Earth-Science Reviews* **155**, 1–27.
- Greenwood SL, Clason CC, Nyberg J, Jakobsson M and Holmlund P (2017) The Bothnian Sea ice stream: early Holocene retreat dynamics of the south-central Fennoscandian ice sheet. *Boreas* **46**(2), 346–362.
- Greenwood SL, Simkins LM, Winsborrow MC and Bjarnadóttir LR (2021) Exceptions to bed-controlled ice sheet flow and retreat from glaciated continental margins worldwide. *Science Advances* **7**(3), eabb6291.

- Gustafson CD and 6 others** (2022) A dynamic saline groundwater system mapped beneath an Antarctic ice stream. *Science* **376**(6593), 640–644.
- Hermanowski P and Piotrowski JA** (2019) Groundwater flow under a paleo-ice stream of the Scandinavian ice sheet and its implications for the formation of Stargard drumlin field, NW Poland. *Journal of Geophysical Research: Earth Surface* **124**(7), 1720–1741.
- Hewitt I** (2011) Modelling distributed and channelized subglacial drainage: the spacing of channels. *Journal of Glaciology* **57**, 302–314.
- Hewitt IJ and Creyts TT** (2019) A model for the formation of eskers. *Geophysical Research Letters* **46**(12), 6673–6680.
- Hoffman MJ and 6 others** (2016) Greenland subglacial drainage evolution regulated by weakly connected regions of the bed. *Nature Communications* **7**(1), 1–12.
- Horgan HJ and 6 others** (2013) Estuaries beneath ice sheets. *Geology* **41**(11), 1159–1162.
- Kamb B** (1987) Glacier surge mechanism based on linked cavity configuration of the basal water conduit system. *Journal of Geophysical Research: Solid Earth* **92**(B9), 9083–9100.
- Kapitsa AP, Ridley JK, de Robin GQ, Siegert MJ and Zotikov IA** (1996) A large deep freshwater lake beneath the ice of central East Antarctica. *Nature* **381**(6584), 684–686.
- Kehew AE, Piotrowski JA and Jørgensen F** (2012) Tunnel valleys: concepts and controversies – a review. *Earth-Science Reviews* **113**(1–2), 33–58.
- Kirkham JD and 6 others** (2020) Morphometry of bedrock meltwater channels on Antarctic inner continental shelves: implications for channel development and subglacial hydrology. *Geomorphology* **370**, 107369.
- Kirkham JD and 6 others** (2021) Tunnel valley infill and genesis revealed by high-resolution 3-D seismic data. *Geology* **49**(12), 1516–1520.
- Kirkham JD and 6 others** (2022) Tunnel valley formation beneath deglaciating mid-latitude ice sheets: observations and modelling. *Quaternary Science Reviews*, 107680.
- Kyrke-Smith TM, Katz RF and Fowler AC** (2014) Subglacial hydrology and the formation of ice streams. *Proceedings of the Royal Society A: Mathematical Physical and Engineering Sciences* **470**(2161), 20130494.
- Larter RD and 6 others** (2019) Subglacial hydrological control on flow of an Antarctic Peninsula palaeo-ice stream. *The Cryosphere* **13**(6), 1583–1596.
- Le Brocq AM and 6 others** (2013) Evidence from ice shelves for channelized meltwater flow beneath the Antarctic ice sheet. *Nature Geoscience* **6**(11), 945–948.
- Lee JR, Wakefield OJ, Phillips E and Hughes L** (2015) Sedimentary and structural evolution of a relict subglacial to subaerial drainage system and its hydrogeological implications: an example from Anglesey, north Wales, UK. *Quaternary Science Reviews* **109**, 88–110.
- Lepp AP and 6 others** (2022) Sedimentary signatures of persistent subglacial meltwater drainage from Thwaites Glacier, Antarctica. *Frontiers of Earth Science* **10**.
- Lewington EL, Livingstone SJ, Clark CD, Sole AJ and Storrar RD** (2020) A model for interaction between conduits and surrounding hydraulically connected distributed drainage based on geomorphological evidence from Keewatin, Canada. *The Cryosphere* **14**(9), 2949–2976.
- Livingstone SJ and 6 others** (2020) A quasi-annual record of time-transgressive esker formation: implications for ice-sheet reconstruction and subglacial hydrology. *The Cryosphere* **14**(6), 1989–2004.
- Mäkinen J** (2003) Time-transgressive deposits of repeated depositional sequences within interlobate glaciofluvial (esker) sediments in Köyliö, SW Finland. *Sedimentology* **50**(2), 327–360.
- Mäkinen J, Kajuutti K, Palmu JP, Ojala A and Ahokangas E** (2017) Triangular-shaped landforms reveal subglacial drainage routes in SW Finland. *Quaternary Science Reviews* **164**, 37–53.
- Minchew BM and Meyer CR** (2020) Dilation of subglacial sediment governs incipient surge motion in glaciers with deformable beds. *Proceedings of the Royal Society A* **476**(2238), 20200033.
- Ojala AE and 6 others** (2019) Ice-sheet scale distribution and morphometry of triangular-shaped hummocks (murtoos): a subglacial landform produced during rapid retreat of the Scandinavian ice sheet. *Annals of Glaciology* **60**(80), 115–126.
- Ojala AE, Mäkinen J, Kajuutti K, Ahokangas E and Palmu JP** (2022) Subglacial evolution from distributed to channelized drainage: evidence from the Lake Murtoo area in SW Finland. *Earth Surface Processes and Landforms* **47**(12), 2877–2896.
- O'Regan M and 6 others** (2021) The Holocene dynamics of Ryder Glacier and ice tongue in north Greenland. *The Cryosphere* **15**(8), 4073–4097.
- Peterson G, Johnson MD, Dahlgren S, Pässe T and Alexanderson H** (2018) Genesis of hummocks found in tunnel valleys: an example from Hörda, Southern Sweden. *GFF* **140**(2), 189–201.
- Rada C and Schoof C** (2018) Channelized, distributed, and disconnected: subglacial drainage under a valley glacier in the Yukon. *The Cryosphere* **12**(8), 2609–2636.
- Röthlisberger H** (1972) Water pressure in intra- and subglacial channels. *Journal of Glaciology* **11**(62), 177–203.
- Rüther DC and 6 others** (2012) Pattern and timing of the northwestern Barents sea ice sheet deglaciation and indications of episodic Holocene deposition. *Boreas* **41**(3), 494–512.
- Schild KM, Hawley RL and Morriss BF** (2016) Subglacial hydrology at Rink Isbræ, West Greenland inferred from sediment plume appearance. *Annals of Glaciology* **57**(72), 118–127.
- Schoof C** (2010) Ice-sheet acceleration driven by melt supply variability. *Nature* **468**(7325), 803–806.
- Schroeder DM, Blankenship DD and Young DA** (2013) Evidence for a water system transition beneath Thwaites Glacier, West Antarctica. *Proceedings of the National Academy of Sciences* **110**(30), 12225–12228.
- Siegfried MR, Fricker HA, Carter SP and Tulaczyk S** (2016) Episodic ice velocity fluctuations triggered by a subglacial flood in West Antarctica. *Geophysical Research Letters* **43**(6), 2640–2648.
- Simkins LM and 6 others** (2017) Anatomy of a meltwater drainage system beneath the ancestral East Antarctic ice sheet. *Nature Geoscience* **10**(9), 691–697.
- Simkins LM and 5 others** (2021) Topographic controls on channelized meltwater in the subglacial environment. *Geophysical Research Letters* **48**(20), e2021GL094678.
- Stevens D and 5 others** (2022) Effects of basal topography and ice-sheet surface slope in a subglacial glaciofluvial deposition model. *Journal of Glaciology* **69**(274), 1–13.
- Storrar RD, Stokes CR and Evans DJ** (2014) Morphometry and pattern of a large sample (>20 000) of Canadian eskers and implications for subglacial drainage beneath ice sheets. *Quaternary Science Reviews* **105**, 1–25.
- Vérité J and 6 others** (2022) Formation of murtoos by repeated flooding of ribbed bedforms along subglacial meltwater corridors. *Geomorphology* **408**, 108248.
- Walder JS** (1986) Hydraulics of subglacial cavities. *Journal of Glaciology* **32**(112), 439–445.
- Walder J and Hallet B** (1979) Geometry of former subglacial water channels and cavities. *Journal of Glaciology* **23**, 335–346.
- Witus AE and 6 others** (2014) Meltwater intensive glacial retreat in polar environments and investigation of associated sediments: example from Pine Island Bay, West Antarctica. *Quaternary Science Reviews* **85**, 99–118.
- Zoet LK, Muto A, Rawling III JE and Attig JW** (2019) The effects of tunnel channel formation on the Green Bay lobe, Wisconsin, USA. *Geomorphology* **324**, 36–47.