

Correspondence

Epidemiol. Infect. (2012).
doi:10.1017/S0950268811001117
First published online 19 August 2011

Risk factors for listeriosis in Australia

To the Editor

I found the paper ‘A national case-control study of risk factors for Listeriosis in Australia’ [1] most interesting. While at the time of the paper’s submission the Food Standards Australia New Zealand (FSANZ) Listeria brochure was only in English, FSANZ has since produced a Listeria and pregnancy video that has been translated into Arabic, Cantonese, Mandarin, Korean, Thai, Vietnamese, Khmer (Cambodian) and Maori. We chose video as a medium, rather than written translations, as we considered that Australian women from non-English-speaking backgrounds may not necessarily be literate in their own language. The video and translations are

available on our website [2]. We may do more translations in future.

Declaration of Interest

None.

References

1. Dalton CB, *et al.* A national case-control study of risk factors for listeriosis in Australia. *Epidemiology and Infection* 2011; **139**: 437–445.
2. Food Standards Australia New Zealand (FSANZ). Translated material, 2010 (<http://www.foodstandards.gov.au/scienceandeducation/translatedmaterial/>).

PAUL BRENT

Chief Scientist Food Standards Australia New Zealand,
55 Blackall St, Barton ACT 2600, 02 6271 2222

(Email: Paul.brent@foodstandards.gov.au)