

modified palliative care screening tool, volunteer ED physicians were asked to identify adult patients with end-stage, chronic conditions including cancer, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), heart failure (HF), cirrhosis, dementia and/or progressive central nervous system (PCNS) disease. Demographic data were collected from these tools and data regarding patients' visits in both the 6 months prior to and 30 days following their index visits were collected from the ED Information System. Bivariate analyses were completed using Student's *t* and chi-square test. **Results:** A total of 663 patients with end-stage illness were identified; 338 (51%) were female and the median age was 76 (IQR: 63, 85). Cancer was the most common presentation (41%), followed by dementia (23%), COPD (16%), HF (9%), CKD (9%), PCNS disease (9%) and cirrhosis (7%). These patients made a total of 1277 visits in the 6 months prior to and 288 in the 30 days following the index visit. Patients presenting to the EDs with cancer ($p = 0.001$), cirrhosis ($p = 0.005$) and CKD ($p = 0.03$) were more likely to visit an ED in the 6 months prior to their index visit. In contrast, patients presenting with dementia ($p < 0.0001$) and PCNS disease ($p = 0.02$) were significantly less likely to present to an ED in the 6 months prior to their index visit. Patients presenting with cirrhosis or CKD had the highest average number of ED visits in the 6 months prior to their index visit (cirrhosis: 4.59 visits, SD: 3.8, $p < 0.0001$; CKD: 4.39 visits, SD: 3.8, $p = 0.0001$). Of these patients, those presenting with end-stage cirrhosis were significantly more likely to make a return visit to an ED within 30 days after their index visit ($p = 0.014$). **Conclusion:** Cancer is the most common condition for patients with end-stage, chronic illnesses in these EDs. Those presenting with cirrhosis or CKD are at a significantly higher risk of repeat visits to the EDs. This study has identified potential deficits in care and can serve as a baseline for future intervention studies.

Keywords: emergency department, palliative care

MP09

Critical care skills training day for emergency medicine residents: A curriculum in evolution

J. Riggs, BA, BSc, S. Gray, MD, MPH, M. McGowan, MHK, A. Petrosniak, MD, MSc, Schulich School of Medicine, Western University; Department of Emergency Medicine, St. Michael's Hospital, Toronto, ON

Introduction: Emergency medicine (EM) residents are expected become proficient in a number of rarely performed, high risk procedures. We developed Critical Care Skills Training Day for senior FRCP and CCFP EM residents at a single university program to fill a gap in resident confidence with these procedures. The day applies principles of deliberate practice with focused feedback using simulation-based training for several rarely performed procedures including thoracotomy, fibre-optic intubation, pericardiocentesis, resuscitative hysterotomy and central line insertion. The objectives of this work was to improve the residents' scores of self-perceived comfort independently performing these procedures by completion of the training day. **Methods:** Clinician educators, residency program directors and simulation specialists designed and taught the curriculum. We used pre- and post-training day surveys blending Likert, multiple choice and free text comments to measure comfort performing each procedure, overall satisfaction and usefulness of this training. Descriptive statistics were used to analyze results. Pre-post differences were assessed using paired sample *T*-tests. Comments and themes from course evaluations were used to make yearly iterative changes.

Results: A total of 95 residents completed the curriculum between 2016-2018. 89 completed evaluations (93%). Residents reported significant ($p < 0.05$) improvement in comfort independently performing fibre optic intubation, thoracotomy and central line insertion. The day was rated very highly, 9.4/10 (SD, 0.72), over 3 years. Feedback was positive with participants identifying opportunities for repeated practice, feedback from instructors and practical tips to improve performance as valuable aspects. Iterative changes were made yearly in response to resident feedback including introduction of new procedures, incorporating skills into sim-based cases, and different training models for skill training. **Conclusion:** Critical Care Skills Training Day for EM residents was created using the principle of deliberate practice to fill a perceived gap in resident training. Residents who completed the annual curriculum showed a marked increase in comfort independently performing several of the procedures. Ongoing challenges include the length of the day, economies of scale, and training models available for the rare procedures. Future directions include the integration of longitudinal objective performance evaluations to align with the competency by design curriculum.

Keywords: deliberate practice, procedural skills, simulation

MP10

Does early intervention improve outcomes for patients with acute ureteral colic?

G. Innes, MD, MSc, A. McRae, MD, PhD, E. Grafstein, MD, J. Andruchow, MD, MSc, M. Law, PhD, F. Scheuermeyer, MD, MSc, University of Calgary, Calgary, AB

Introduction: The optimal initial management approach for ureteral colic is unclear. Guidelines recommend spontaneous passage for most patients, but early stone intervention may rapidly terminate acute episodes. We compared 60-day treatment failure rates in matched patients undergoing early intervention versus spontaneous passage. **Methods:** We used administrative data and structured chart review to study all emergency department (ED) patients at nine Canadian hospitals who had an index ureteral colic visit and a computed tomography (CT) confirmed 2.0-9.9 mm stone during 2014. Using Cox Proportional Hazards models, we assessed 60-day treatment failure, defined as hospitalization or rescue intervention, in patients undergoing early intervention compared to propensity-score matched controls undergoing trial of spontaneous passage. **Results:** From 3,081 eligible patients, mean age 51 years and 70% male, we matched 577 patients in each group (total 1154). Control and intervention cohorts were balanced on all parameters and propensity scores, which reflect the conditional probability a patient would undergo early intervention, were similarly distributed. In the time to event analysis, 21.8% in both groups experienced the composite primary outcome of treatment failure (difference = 0%; 95% CI, -4.8 to 4.8%). Early intervention patients required more ED revisits (36.1% v. 25.5%; difference 10.6%; 95% CI 5.3 to 15.9%) and more 60-day hospitalizations (20.1% v. 12.8%). The strongest predictors of adverse outcome were stone size, proximal or middle stone location, and ED length of stay. **Conclusion:** If applied broadly to patients with 2.0-9.9mm ureteral stones, an early interventional approach was associated with similar rates of treatment failure, but more hospitalizations and emergency revisits. Research clarifying subgroups most likely to benefit will facilitate better targeting of early intervention, potentially reducing patient morbidity and improving system utilization.

Keywords: intervention, outcomes, renal colic